

THE SPATIAL EXTENT OF (U)LIRGs IN THE MID-INFRARED. II. FEATURE EMISSION

T. DÍAZ-SANTOS^{1,†}, V. CHARMANDARIS^{1,2}, L. ARMUS³, S. STIERWALT³, S. HAAN³, J. M. MAZZARELLA⁴, J. H. HOWELL³, S. VEILLEUX⁵, E. J. MURPHY⁶, A. O. PETRIC³, P. APPLETON⁷, A. S. EVANS^{8,9}, D. B. SANDERS¹⁰, AND J. A. SURACE³

Draft version August 1, 2011

ABSTRACT

We present results from the second part of our analysis of the extended mid-infrared (MIR) emission of the Great Observatories All-Sky LIRG Survey (GOALS) sample based on $5 - 14 \mu\text{m}$ low-resolution spectra obtained with the Infrared Spectrograph on *Spitzer*. We calculate the fraction of extended emission as a function of wavelength for all galaxies in the sample, FEE_λ , defined as the fraction of the emission that originates outside of the unresolved central component of a source, and spatially separate the MIR spectrum of a galaxy into its nuclear and extended components.

We find that the $[\text{Ne II}]12.81 \mu\text{m}$ emission line is as compact as the hot dust MIR continuum, while the polycyclic aromatic hydrocarbon (PAH) emission is more extended. In addition, the 6.2 and $7.7 \mu\text{m}$ PAH emission is more compact than that of the $11.3 \mu\text{m}$ PAH, which is consistent with the formers being enhanced in a more ionized medium. The presence of an AGN or a powerful nuclear starburst increases the compactness and the luminosity surface density of the hot dust MIR continuum, but has a negligible effect on the spatial extent of the PAH emission on kpc-scales. Furthermore, it appears that both processes, AGN and/or nuclear starburst, are indistinguishable in terms of how they modify the integrated PAH-to-continuum ratio of the FEE in (U)LIRGs. Globally, the $5 - 14 \mu\text{m}$ spectra of the extended emission component are homogeneous for all galaxies in the GOALS sample. This suggests that, independently of the spatial distribution of the various MIR features, the physical properties of star formation occurring at distances farther than 1.5 kpc from the nuclei of (U)LIRGs are very similar, resembling local star-forming galaxies with $L_{\text{IR}} < 10^{11} L_\odot$, as well as star formation-dominated ULIRGs at $z \sim 2$. In contrast, the MIR spectra of the nuclear component of local ULIRGs and LIRGs are very diverse. These results imply that the observed variety of the integrated MIR properties of local (U)LIRGs arise, on average, only from the processes that are taking place in their cores.

Subject headings: infrared: galaxies — galaxies: evolution — galaxies: interactions — galaxies: starburst — galaxies: active

1. INTRODUCTION

The mid-infrared (MIR) spectra of galaxies provide key information that allow us to characterize the global properties of their star formation, as well as to develop continuum and emission line diagnostics in order to infer whether they harbor an active galactic nucleus (AGN), with less confusion due to dust extinction compared to observations at optical wavelengths (Lutz et al.

1998b; Genzel et al. 1998; Charmandaris et al. 2004; Brandl et al. 2006; Smith et al. 2007; Armus et al. 2007; Veilleux et al. 2009; Wu et al. 2009; Sales et al. 2010; Petric et al. 2011). However, when studying the integrated emission of galaxies, the spatial distributions of the sources contributing to their observed infrared (IR) spectral energy distributions, such as H II regions, photo-dissociation regions (PDRs), cold molecular clouds, and/or non-thermal emission sources, are averaged together. That is, all the intrinsic properties of the different phases of the inter-stellar medium (ISM) are mixed, despite the fact that they can be substantially different (Laurent et al. 2000; Smith et al. 2004). Therefore, even a rough decomposition between the nuclear and disk components of a galaxy can yield valuable information on where the IR luminosity originates, as well as details on the processes associated to the different emission components.

The access to this spatial information is even more vital in the study of luminous and ultra-luminous infrared galaxies, (U)LIRGs, for which it is known that the bulk of their energy production is generated within their central regions and emitted in the IR on scales of a few kpc. This class of galaxies, although not very numerous in the nearby Universe (Sanders & Mirabel 1996), are responsible for the bulk of the obscured star formation at $z \geq 1$ (Le Floc'h et al. 2005; Pérez-González et al. 2005;

¹ University of Crete, Department of Physics, GR-71003, Heraklion, Greece: tanio@physics.uoc.gr

[†] Now at the Spitzer Science Center, California Institute of Technology, MS 220-6, Pasadena, CA 91125

² IESL/FORTH - Department of Physics and Institute of Theoretical & Computational Physics, 71003, Heraklion, Greece and Chercheur Associé, Observatoire de Paris, F-75014, Paris, France

³ Spitzer Science Center, California Institute of Technology, MS 220-6, Pasadena, CA 91125

⁴ Infrared Processing & Analysis Center, MS 100-22, California Institute of Technology, Pasadena, CA 91125

⁵ Department of Physics, University of Oregon, Eugene, OR 97403

⁶ Observatories of the Carnegie Institution for Science, 813 Santa Barbara Street, Pasadena, CA 91101

⁷ NASA Herschel Science Center, IPAC, MS 100-22, California Institute of Technology, Pasadena, CA 91125

⁸ Department of Astronomy, University of Virginia, P.O. Box 400325, Charlottesville, VA 22904

⁹ National Radio Astronomy Observatory, 520 Edgemont Road, Charlottesville, VA 22903

¹⁰ Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822

Caputi et al. 2007; Magnelli et al. 2011; Murphy et al. 2011). Thus, the study of the spectra of local (U)LIRGs, where the angular resolution achieved by space-born observatories such as *Spitzer* is sufficient to spatially-resolve their components, is essential for interpreting the internal mechanisms that govern their IR emission and for helping to understand the properties of high redshift (U)LIRGs.

Several MIR spectroscopic studies have already been carried out using ground-based telescopes which provide, for the closest and brightest sources, spatial resolution of a few tens of pc (Soifer et al. 2002; Soifer et al. 2003; Díaz-Santos et al. 2010a). These works have demonstrated that the MIR continuum emission, ionization lines, and polycyclic aromatic hydrocarbon (PAH) features arise from different regions in (U)LIRGs, and that the sources of emission are arranged in more compact configurations than in galaxies with lower IR luminosities. This is consistent with the merger-induced nature of most local (U)LIRGs (Veilleux et al. 2002; Armus et al. 2009) and the funneling of large quantities of gas and dust mass towards their nuclei during the interaction. In addition, MIR spectral maps using the IRS instrument on-board *Spitzer* have been obtained for 15 local LIRGs revealing an increase of the electron density in their nuclei (Pereira-Santaella et al. 2010).

Despite the small number of galaxies studied in these works, some general conclusions about the properties of (U)LIRGs as a whole could be derived. Yet, these trends are confounded by systematics caused by intrinsic variations of processes that dominate at sub-kpc scales. Studying these internal processes is therefore essential if we wish to completely understand the global trends. The Great Observatories All-sky LIRG Survey (GOALS; Armus et al. 2009), which consist of 291 galaxies (202 systems), is the ideal local LIRG sample for this task because it is large enough to allow us to break it up into sub-samples based, for example, on their interaction stage, or the AGN contribution to their MIR emission (Petric et al. 2011). Each sub-sample has enough galaxies in order to perform a robust statistical analysis and examine how the various MIR properties vary from one sub-sample to another and within a given sub-sample. Such an analysis was performed by Díaz-Santos et al. (2010b), hereafter Paper I, who analyzed the compactness of the MIR continuum emission at $13.2\mu\text{m}$ of (U)LIRGs in the GOALS sample and discussed its correlation to several physical properties of the systems such as their L_{IR} , MIR AGN fraction, and far infrared (FIR) colors.

The wealth of information available in the $5 - 14\mu\text{m}$ spectral range allow us to further expand our analysis to several other key MIR features. In this second paper we explore how the compactness of the emission due to polycyclic aromatic hydrocarbon (PAH) and atomic lines compares to that of the warm dust continuum and investigate the origin of the observed differences. An analysis regarding the nuclear PAH emission ratios among the GOALS sample will be further addressed in a following study by Stierwalt et al. (2011, in prep.). The paper is structured as follows: in Section 2 we briefly remind the reader of the datasets used and the methodology of our analysis which was described in detail in Paper I. Our results are reported in Section 3, where we present average MIR spectra of the extended and nuclear emission of the

three main types of sources we found in Paper I. We also explore the role of dust extinction as well as the presence of an active galactic nucleus (AGN) in the observed spatial profiles of the various MIR spectral features. Finally, in Section 4 we present our conclusions.

2. OBSERVATIONS AND DATA ANALYSIS

2.1. The sample

The sample we analyze is the Great Observatories All-Sky LIRG Survey (GOALS; Armus et al. 2009). GOALS comprises a complete, flux-limited sample of galaxies in the local Universe drawn from the Revised Bright Galaxy Sample (RBGS, Sanders et al. 2003) selected to be systems in the (U)LIRG luminosity classes. Armus et al. (2009) describe in detail how the sample was selected as well its global characteristics. In addition, using a number of MIR diagnostics, Petric et al. (2011) estimate the AGN contribution to the MIR luminosity of the systems and, based on their apparent morphology, classify each galaxy into a stage of interaction, ranging from isolated systems to advanced mergers. In Howell et al. (2010) the relation between the UV and MIR emission of the galaxies in the sample is investigated, while Haan et al. (2011) present a thorough analysis of the nuclear structure of the galaxy sample using high spatial resolution NIR and optical images obtained with the *Hubble Space Telescope*. We refer the reader to these papers since we will rely on their findings for the interpretation of our results. Out of the 291 galaxies (202 systems) included in the GOALS sample, a total of 221 are used for this study, 200 LIRGs and 21 ULIRGs. All of them have low-resolution spectroscopic *Spitzer*/IRS observations in the $5 - 14\mu\text{m}$ rest-frame wavelength range. The angular resolution of IRS ($\sim 3''.6$ at $13.2\mu\text{m}$) allows us to separate regions of physical scales ranging from 0.22 kpc at the distance of the closest LIRG ($\sim 12\text{Mpc}$), to 6.1 kpc at $\sim 340\text{Mpc}$ where the farthest ULIRG of the sample is located. The spatial resolution at the median distance of our galaxy sample, 91 Mpc, is 1.7 kpc (also at $13.2\mu\text{m}$). A table with the main characteristics of the galaxies such as their distance, L_{IR} , or FIR colors can be found in Paper I. We also analyzed separately a compilation of ULIRGs from the works of Imanishi et al. (2007); Imanishi (2009); Imanishi et al. (2010), but these sources cannot be included in our present study because they are too distant and, as a consequence, they are unresolved at all wavelengths observed with the IRS.

2.2. Analysis

Our analysis is based on the calculation of the fraction of extended emission of galaxies as a function of wavelength, FEE_λ , which is defined as the fraction of emission in a galaxy that does *not* arise from its spatially unresolved central component. Detailed information on the calculation of the FEE_λ functions can be found in Paper I. For practical purposes, we remind the reader that its formal definition is:

$$\text{FEE}_\lambda = \frac{\text{EE}_\lambda}{\text{E}_\lambda(\text{total})} \quad (1)$$

where FEE_λ , EE_λ and $\text{E}_\lambda(\text{total})$ are the fraction extended emission (ranging from zero to unity), the extended emission, and the total emission of a galaxy




Figure 1. The FEE_λ function, smoothed with a 4-pixel box to reduce noise, are plotted in blue for 3 galaxies that serve as examples of the main types identified in the sample. The Spitzer/IRS $\sim 5\text{--}14\mu\text{m}$ spectrum of each galaxy, scaled to arbitrary units, is also plotted as a pink dashed line for reference. Left panel: NGC 3110 with constant/featureless FEE_λ . Middle panel: NGC 1365 displaying a PAH and line extended emission. Right panel: MCG+08-11-002 with silicate-extended emission.

within the IRS slit at each wavelength respectively. Essentially, the FEE_λ of a source can be considered as the complementary part of its compactness, which can be defined as $1 - FEE_\lambda$. We also define the core size of a galaxy as the full width half-maximum (FWHM) of a Gaussian fitted to the spatial profile of its nuclear emission along the *Spitzer*/IRS slit at a given wavelength. We note that while the core size represents how extended the nuclear emission is, the FEE also accounts for low surface brightness emission that is more extended than the core of the source (beyond the wings of the central Gaussian). In Table 2 (available in electronic format) we provide the FEE_λ measurements at the wavelengths of the MIR spectral features considered in this work.

3. RESULTS AND DISCUSSION

In Paper I we grouped the GOALS galaxies into 3 types of FEE_λ functions: constant/featureless, PAH- and line-extended, and silicate-extended. We excluded 8 sources that appear unresolved, thus having an $FEE_{5-14\mu\text{m}} \simeq 0$, and 11 that could not be classified in any of the three main FEE_λ types. However, for completeness, in the present paper we do take them into account. Figure 1 shows examples of the FEE_λ function types.

3.1. The FEE of Continuum, PAHs, and Emission Lines

In Figure 2 we examine how the FEE of several MIR spectral features, averaged at the wavelength where they peak $\pm 0.1\mu\text{m}$, vary as a function of the FEE of the continuum emission at $13.2\mu\text{m}$, which was studied in Paper I. From left to right and top to bottom we display: the $11.3\mu\text{m}$ PAH FEE as a function of the FEE at the peak of the $9.7\mu\text{m}$ silicate feature, and the $6.6\mu\text{m}$ continuum FEE, [Ne II] $12.81\mu\text{m}$ emission line (+ $12.5\mu\text{m}$ PAH) FEE, $11.3\mu\text{m}$ PAH FEE, $7.7\mu\text{m}$ PAH FEE, and $6.2\mu\text{m}$ PAH FEE as a function of the $13.2\mu\text{m}$ continuum FEE.

Figure 2a shows, in a quantitative way, the visual classification of the FEE_λ functions of galaxies performed in Paper I, which we used for separating sources with characteristic FEE_λ in three main types/categories (Figure 1). Within the uncertainties, (U)LIRGs showing a

constant/featureless FEE_λ (blue diamonds) are located on the one-one relation (black dashed line). Sources presenting an excess of extended PAH emission with respect to that of the continuum (green stars) are above the line, while galaxies with the FEE_λ peaking at $9.7\mu\text{m}$ (red circles) are generally located below. We refer the reader to Paper I for more details about this classification.

The principal reason we plot the $6.6\mu\text{m}$ continuum in Figure 2b is to verify that the loss of angular resolution as a function of wavelength that affects the $5 - 14\mu\text{m}$ SL spectra is not biasing our measurements of the FEE_λ , such that it decreases as wavelength increases. We can see that this is not the case, as the best fit to the $FEE_{6.6\mu\text{m}}$ and $FEE_{13.2\mu\text{m}}$ data, both representing continuum emission, is compatible, within the uncertainty (see the gray shadow), with the one-to-one relation. This implies that the decrease in angular resolution as the wavelength increases does not affect the trends significantly, at least in terms of the FEE measurements, when enough galaxies are considered.

Figure 2c shows the FEE averaged at the location of the [Ne II] $12.81\mu\text{m}$ emission line as a function of $FEE_{13.2\mu\text{m}}$. Both quantities are well correlated, which is in agreement with previous studies of smaller samples of local LIRGs (Pereira-Santaella et al. 2010), and implies that the physical processes (AGN and/or star formation) that are responsible for the [Ne II] $12.81\mu\text{m}$ line and the MIR continuum emission are equally concentrated and that they are likely the same. On the other hand, the FEE of the $11.3\mu\text{m}$ PAH is larger than that of the continuum or the [Ne II] $12.81\mu\text{m}$ emission in many galaxies (Figure 2d). Moreover, this is independent of the FEE_λ function type and does not only occur for silicate-dominated FEE_λ type galaxies (red circles; see related caveat in Section 3.2, where we discuss the effect of the extinction on the spatial distribution of the MIR emission of spectral features located within the wavelength range of the $9.7\mu\text{m}$ silicate absorption band). This result is in agreement with works based on high spatial resolution *N*-band ($8 - 13\mu\text{m}$) spectroscopy of local LIRGs where it has been found that the spatial distribution of the [Ne II] $12.81\mu\text{m}$ emis-




Figure 2. From left to right and top to bottom: (a) FEE_{11.3μm} (11.3 μm PAH) vs. FEE_{9.7μm} (peak of the silicate absorption feature). Then, FEE_{13.2μm} (continuum) vs.: (b) FEE_{6.6μm} (continuum), (c) FEE_{12.8μm} ([Ne II] 12.81 μm line), (d) FEE_{11.3μm}, (e) FEE_{7.7μm} (7.7 μm PAH), and (f) FEE_{6.2μm} (6.2 μm PAH). The different symbols and colors represent the FEE_λ galaxy types (unresolved: dark grey inverted triangles; constant/featureless: blue diamonds; PAH-dominated: green stars; silicate-dominated: red circles; and unclassified: yellow squares). The black dashed line shows the one-to-one relation, while the black solid line is the linear fit to the data, whose parameters can be found at the bottom right of the plots. The shaded region is the 1σ uncertainty of the fit, while the light gray underlying line is the arbitrarily-scaled histogram of the data distributed in equal bins of FEE_{9.7μm} or FEE_{13.2μm}.

sion and the MIR continuum in star-forming regions follows that of the Pa α emission (Díaz-Santos et al. 2010a), which is in turn associated to young, compact H II regions (e.g., Alonso-Herrero et al. 2002). Meanwhile, the 11.3 μ m PAH emission can be powered not only by H II regions but also by rather “evolved” ($\gtrsim 8$ –10 Myr) stars, which emit fewer ionizing photons. In other words, the 11.3 μ m PAH/[Ne II]12.81 μ m flux ratio depends on the age of the ionizing stellar populations (Díaz-Santos et al. 2010a). This diffuse PAH emission would be located in between the circumnuclear young star-forming regions and also farther from the nucleus of (U)LIRGs (Díaz-Santos et al. 2008; Pereira-Santaella et al. 2010; Bendo et al. 2010). One might think that, because of the limited spectral resolution of the data, the [Ne II]12.81 μ m line could be contaminated by the emission of the 12.5 μ m PAH feature. However, this is not the case, because if the 12.5 μ m PAH were as extended as the 11.3 μ m PAH, or even as the 7.7 or 6.2 μ m PAHs (see below), and the 12.5 μ m PAH was dominating the emission at 12.8 μ m then, for a given value of the $FEE_{13.2\mu\text{m}}$, the $FEE_{12.8\mu\text{m}}$ should appear much more extended and/or with a larger scatter than what is observed in Figure 2c. Instead, one can identify a tight correlation between both FEEs at 12.8 and 13.2 μ m.

Figure 2e and 2f show that the 7.7 and 6.2 μ m PAH emissions are as compact as, or just slightly more extended than, that of the MIR continuum and hence the [Ne II]12.81 μ m emission, but less than the 11.3 μ m PAH. This is in agreement with the notion that the 7.7 and 6.2 μ m emission bands are enhanced for ionized PAHs, while the 11.3 μ m PAH emission is similar for neutral and ionized molecules (e.g., Allamandola et al. 1999; Galliano et al. 2008). Since ionized PAH carriers require harder radiation fields in order to be excited than neutral PAHs (Draine & Li 2001), they will be more tightly associated to younger massive star forming regions which, in turn, are also strong [Ne II]12.81 μ m emitters. Given the nature of LIRGs and ULIRGs, many of which harbor strong nuclear starbursts, this would result in having both the 7.7 and the 6.2 μ m PAHs slightly more extended than the [Ne II]12.81 μ m line but less than the 11.3 μ m PAH.

In Paper I we also examined the variations of the FEE of the continuum at 13.2 μ m as a function of the L_{IR} . Here we also explore how the FEE of PAHs in (U)LIRGs depends on the L_{IR} and compare them with their corresponding $FEE_{13.2\mu\text{m}}$. Figure 3 shows that, despite the large dispersion, the median of the $FEE_{7.7\mu\text{m}}/FEE_{13.2\mu\text{m}}$ ratio (black solid line) slightly increases with L_{IR} . The more IR luminous a galaxy is, the more extended its 11.3 μ m PAH emission is compared with the MIR continuum. This is in agreement with our earlier result in Paper I, where we found that the compactness of the MIR continuum increases for galaxies with $L_{\text{IR}} \gtrsim 10^{11.8} L_{\odot}$. However, the 6.2, 7.7, and 11.3 μ m PAHs do not show such a steep decrease of their FEEs around these IR luminosities. Therefore, it is mainly the MIR continuum that becomes more compact as the L_{IR} of galaxies increases, making the median PAH-to-continuum FEE ratio to increase too. Likewise, the most luminous galaxies are also those having the warmest $\log(f_{60\mu\text{m}}/f_{100\mu\text{m}})$ FIR ratios. This is expected for ULIRGs, which for a long time are known to host nuclear starbursts and AGNs




Figure 3. $FEE_{7.7\mu\text{m}}/FEE_{13.2\mu\text{m}}$ ratio as a function of the L_{IR} of our (U)LIRG sample. The symbols and colors depend on the IRAS $\log(f_{60\mu\text{m}}/f_{100\mu\text{m}})$ FIR ratio of the galaxies. The blue squares and red circles represent cold and warm sources with $\log(f_{60\mu\text{m}}/f_{100\mu\text{m}})$ ratios lower and larger than -0.2 respectively. The blue and red filled diamonds show the median values of the $FEE_{7.7\mu\text{m}}/FEE_{13.2\mu\text{m}}$ ratio for the two groups of sources in equally-distributed IR luminosity bins. The black diamonds represent the same ratio but considering all sources. Galaxies whose core sizes are unresolved at 13.2 μ m are marked with yellow crosses.

and show on average warm colors (Sanders & Mirabel 1996), but these same properties are also displayed by several sources with lower IR luminosities ($10^{11} L_{\odot} \lesssim L_{\text{IR}} \lesssim 10^{12} L_{\odot}$). Indeed, independently of their L_{IR} , the warmest galaxies, those with $\log(f_{60\mu\text{m}}/f_{100\mu\text{m}}) > -0.2$ (identified by red circles in Figure 3), are those driving the trend and accounting for the dispersion seen in the data. As we will discuss later, this is in agreement with their compact nature and the general scenario we will propose in Section 3.4.

3.2. The Role of Dust Extinction

We also investigated the specific distribution of (U)LIRGs of different FEE_{λ} types in Figure 2a–e. As we mentioned above, Figure 2c shows that the $FEE_{11.3\mu\text{m}}$ is larger than the $FEE_{13.2\mu\text{m}}$ in many (U)LIRGs. However, for some of them this may be an extinction effect.

Even in the MIR, dust obscuration must be taken into account in (U)LIRGs since their nuclear extinction is much larger than that of their extended regions (e.g., García-Marín et al. 2009). Galaxies with a silicate-dominated FEE_{λ} type are marked as red circles in Figure 2. We observe that for these sources the 11.3 μ m PAH feature, which is located within the 9.7 μ m silicate absorption feature, appears more extended (i.e., have large $FEE_{11.3\mu\text{m}}$ values). This is not because there is more PAH emission in the disks of these galaxies but instead because the emission is suppressed in their nuclei due to the extinction, as we already suggested in Paper I. Therefore, the differential extinction between the unresolved and extended components plays an important role in the “apparent” spatial distribution of the




Figure 4. Histogram of the $FEE_{7.7\mu m}/FEE_{13.2\mu m}$ ratio of our galaxies grouped as a function of the AGN contribution to their MIR continuum. A total of 205 (U)LIRGs for which we have measurements of their $FEE_{7.7\mu m}/FEE_{13.2\mu m}$ ratios and AGN-fractions from Petric et al. (2011) were used. The solid lines represent the median value of each distribution, while the dotted lines show the $\pm 1\sigma$ uncertainty with respect to the medians. A Kolmogorov-Smirnov test indicates clearly ($p \sim 10^{-6}$) that the distributions are not drawn from same parent population.

emission of the different MIR features in these galaxies. Sources with extreme differences between the extinction in their nuclei and extended regions (large $\Delta S_{Si\,9.7\mu m}$, see Section 3.5) will show an artificial excess in their $FEE_{11.3\mu m}/FEE_{13.2\mu m}$ ratio (see implications in next Sections). Indeed, this also applies in general to all features within the $\sim 8-12.5\mu m$ wavelength range affected by the $9.7\mu m$ silicate feature, such as the $8.6\mu m$ PAH and the [SIV] $10.51\mu m$ line. As a consequence, in order to interpret correctly the FEE ratios of those galaxies with large $\Delta S_{Si\,9.7\mu m}$, a careful decomposition of their spectra in different dust components and feature emissions must be done to properly account for extinction effects (Stierwalt et al. 2011, in prep.).

Nevertheless, not all (U)LIRGs have highly obscured nuclei at $9.7\mu m$. There are several galaxies in our sample, those with a PAH-dominated FEE_λ type, that show high PAH-to-continuum FEE ratios but, unlike galaxies with silicate-dominated FEE_λ types, they do not display large differences between the extinction of their unresolved and extended components.

3.3. The Influence of an AGN

In Paper I we studied how the FEE of the continuum at $13.2\mu m$ varies with the MIR AGN fraction of galaxies and showed that the MIR continuum becomes more compact for AGN-dominated sources. In order to examine whether the presence of an AGN also influences the distribution of the PAH emission in (U)LIRGs, we plot in Figure 4 the histogram of the ratio of the $7.7\mu m$ PAH FEE to the $13.2\mu m$ continuum FEE, grouping the galaxies as a function of their MIR AGN-fraction. Galaxies in which an AGN dominates the MIR emis-

sion display large PAH-to-continuum FEE ratios. The median $7.7\mu m$ PAH-to-continuum FEE ratio for MIR star formation-dominated galaxies, that is, sources for which Petric et al. (2011) find that the AGN contribution to the MIR emission is less than 50%, is 1.08 ± 0.10 , while for AGN-dominated sources, it is 1.86 ± 0.41 . The corresponding median ratios for the $6.2\mu m$ PAH-to-continuum FEE ratio are 1.04 ± 0.11 and 1.74 ± 0.51 respectively. We note that a much larger difference is seen in the $FEE_{11.3\mu m}/FEE_{13.2\mu m}$ ratio. However, as discussed in Section 3.2, the $FEE_{11.3\mu m}$ is affected by the $9.7\mu m$ silicate absorption feature and, at the same time, AGN-dominated galaxies tend to show large differences between the extinction of their nuclei and extended regions, i.e., large $\Delta S_{Si\,9.7\mu m}$ values. Both effects combined may artificially increase the $11.3\mu m$ PAH-to-continuum FEE ratio leading to a possible misinterpretation of the observed correlation. The 7.7 and $6.2\mu m$ PAH, on the other hand, do not suffer from severe obscuration and therefore the trend and ratios seen in Figure 4 are more reliable.

Although the difference between the median PAH-to-continuum FEE ratios of MIR AGN and star formation-dominated galaxies is at the $\sim 2\sigma$ level, a Kolmogorov-Smirnov test performed between the two samples indicates clearly ($p \sim 10^{-6}$) that the distributions are not drawn from same parent population. This, combined with the fact that the $FEE_{13.2\mu m}$ decreases as the MIR AGN fraction increases (see Paper I) suggest that as the MIR emission of a (U)LIRG begins to be dominated by an AGN, the hot dust continuum becomes more compact while the PAH emitting regions do not. That is, the presence of an AGN modifies the spatial extent of the hot dust continuum but has a negligible effect on the overall, kpc-scale distribution of the PAH emission.

Indeed, this is consistent with the way in which the AGN fraction is calculated in Petric et al. (2011). In this work the contribution of an AGN to the MIR emission of a (U)LIRG is estimated through the so-called Laurent diagram (Laurent et al. 2000; Armus et al. 2007) which, for the GOALS sample, relies mainly on the ratio between the $6.2\mu m$ PAH emission and the $5.5\mu m$ continuum (see Figure 3 in Petric et al. 2011). Therefore, as the compactness of the hot dust continuum increases, so does the surface brightness of the galaxy and, as a consequence, this pseudo-equivalent width (EW) of the $6.2\mu m$ PAH decreases, which implies that the AGN contribution to the MIR is larger and dilutes the PAH emission.

We have also investigated the influence of an AGN depending on the FEE_λ type of the host. For example, there are no galaxies with constant/featureless FEE_λ having a MIR AGN fraction above 0.5. This is an important result since, as we have explained, the galaxies were classified into the different FEE_λ types independently of their MIR spectral characteristics. Thus, it appears that when an AGN dominates the MIR emission it also leaves a “footprint” in the FEE_λ function of a galaxy in the sense that the MIR emission of (U)LIRGs with a constant FEE_λ will never be dominated by an AGN or, in general, appear compact.

We would like to stress here that not only AGN-dominated galaxies show large PAH-to-continuum FEE ratios. Nearly 30% of (U)LIRGs whose MIR emission is




Figure 5. Top: $FEE_{7.7\mu m}/FEE_{13.2\mu m}$ ratio as a function of the *IRAS* $\log(f_{60\mu m}/f_{100\mu m})$ FIR color. The red circles and blue squares mark galaxies whose MIR AGN fraction is ≥ 0.5 and < 0.5 respectively. The small black dots represent sources for which the MIR AGN fraction is not available. The dashed line is defined as the median PAH-to-continuum FEE ratio value $+ 2\sigma$ of star-formation dominated sources (see Figure 4). Bottom: Same plot but for the $FEE_{6.2\mu m}/FEE_{13.2\mu m}$ ratio.

not dominated by an AGN (AGN-fraction < 0.5) display large $FEE_{7.7\mu m}/FEE_{13.2\mu m}$ ratios. The nature of these sources will be addressed in the next Section.

3.4. PAH-to-Continuum FEE Ratios and FIR Emission

In this section we examine whether the PAH-to-continuum FEE ratios are related with the FIR colors of galaxies, that is, whether the differences in the compactness of the PAH and MIR continuum emission in (U)LIRGs are linked to their global dust temperatures traced by the FIR emission. We have already discussed that PAH emission probes time-scales of star formation

larger than the MIR continuum emission, which is tightly associated to young, ionizing stellar populations (e.g., Calzetti et al. 2007; Díaz-Santos et al. 2010a). Likewise, results based on *IRAS* as well as on recent high spatial resolution *Herschel* observations have clearly demonstrated that the $100 - 500 \mu m$ emission of galaxies arises mainly from a diffuse cold dust component that extends smoothly along their disks and is heated by old stars, rather than being concentrated in knots of current star formation which are better traced by dust continuum emission at $< 100 \mu m$ (Sauvage & Thuan 1992; Walterbos & Greenawalt 1996; Bendo et al. 2010; Kramer et al. 2010). Moreover, in a recent *Herschel* study of local spiral galaxies it has been found that their cores show enhanced, warmer $70/160 \mu m$ colors than their disks (Sauvage et al. 2010). Therefore, since in Paper I we showed that the $FEE_{13.2\mu m}$ decreases when (U)LIRGs become warmer, i.e., when their *IRAS* $\log(f_{60\mu m}/f_{100\mu m})$ FIR color increases, we would also expect to find a trend between the PAH-to-continuum FEE ratios and the FIR color of galaxies.

Figure 5 shows that this is actually the case. Although the trend for the $6.2 \mu m$ PAH-to-continuum FEE ratio is less significant, it is clear that the $7.7 \mu m$ PAH-to-continuum FEE ratio and its dispersion increase as galaxies become warmer. Moreover, practically all (90%) MIR star formation-dominated sources with large PAH-to-continuum FEE ratios (sources above the dashed lines, which are defined as the medians $+ 2\sigma$ of the distributions shown in Figure 4) have $\log(f_{60\mu m}/f_{100\mu m}) \gtrsim -0.2$. A plausible explanation for this is the following: as the nuclear starburst activity in a (U)LIRG increasingly dominates its integrated IR emission, the dust in the central regions would reach higher temperatures. This is because more and more of the atoms and molecules of the nuclear gas would be ionized and dissociated by the high energy photons of the stars formed in the H II regions of their proximity, causing the ratio between the volume of the spherical shells of the PDRs, from where most of the PAH emission originate, and the volume of the dust emitting region to decrease. That is, a larger fraction of the ionizing photons would be absorbed by dust close to the stars increasing its temperature, and would not reach the PDRs. As a result, the MIR continuum emission due to this hot dust will be more compact (having a higher luminosity surface density), and the global galaxy colors will become warmer.

We could then consider that the “normal” behavior of a galaxy is to show the same spatial extent of the PAH and MIR continuum emission probably as a consequence of a sufficient mixing of its gas and stars. It is only when a strong, compact emitter starts to dominate the nuclear MIR continuum that we see a deviation (i.e., PAH-to-continuum FEE ratios above the dashed lines in Figures 4 and 5). This is the case for 80% of AGN-dominated GOALS galaxies and 60% of warm sources with $\log(f_{60\mu m}/f_{100\mu m}) > -0.2$, excluding AGN-dominated galaxies. In these systems, the MIR continuum is more compact than the PAH emission, which is still as extended as in the “normal” population since it arises mostly from a diffuse component (the disk of the galaxies). Therefore, a more appropriate interpretation would not be that there is an excess of PAH emission or that the PAHs are more extended in absolute

terms in these galaxies. It is the MIR continuum that is more compact due to the enhancement caused by the nuclear power-source that increases the PAH-to-continuum FEE ratio. It is important to stress that these compact galaxy cores are not only observed in ULIRGs (see, e.g., Charmandaris et al. 2002); we also see them in LIRGs (see also Figure 3). In addition, Figures 4 and 5 also suggests that irrespective of whether the nuclear MIR emission of (U)LIRGs is dominated by an AGN or a nuclear starburst, their impact on the compactness of the MIR continuum and PAH emission is the same. In other words, both processes are indistinguishable in terms of the relative spatial extent of the PAH and MIR continuum emission.

We note that the MIR emission of 27% of the warm compact sources, those with $\log(f_{60\mu\text{m}}/f_{100\mu\text{m}}) > -0.2$ and above the dashed lines in the Figures, is dominated by an AGN (MIR AGN fraction ≥ 0.5). It is known that AGN and star formation dominates the emission mainly at different wavelengths, i.e., the FIR colors of these galaxies are not affected by the existence of an AGN but they are associated principally to their star formation processes. This implies that in these (U)LIRGs, nuclear compact star formation, traced by their warm FIR colors, must be coexisting with MIR-detected AGN at the same time (see, e.g., Schweitzer et al. 2006; Netzer et al. 2007; Veilleux et al. 2009), both contributing to the large PAH-to-continuum FEE ratios they show.

3.5. Averaged Spectra of Unresolved and Extended Components

The technique we use to calculate the FEE_λ from the 2 dimensional *Spitzer*/IRS images allows us to derive not only the spatial extent of specific spectral features but also to separate the MIR spectrum of the extended emission component of galaxies from that of the unresolved component in a simple manner. This can be done just by multiplying their total, flux-calibrated *Spitzer*/IRS spectrum calculated by Petric et al. (2011), by their FEE_λ and $1-\text{FEE}_\lambda$ functions respectively. More technical details are available in the Appendix of Paper I.

Using this method, we display together in Figure 6 the averaged spectra of the different FEE_λ types for the unresolved and extended components of (U)LIRGs. The component averages were weighted by the mean FEE_λ and $1-\text{FEE}_\lambda$ of each galaxy, respectively. This was done in order to give proportionally more importance in the calculation of the extended component averages to (U)LIRGs showing larger values of FEE and, on the other hand, more importance to (U)LIRGs showing lower values of FEE when the unresolved component averages are calculated. In any case, we note that the results are the same irrespective of this weighting.

Strikingly, we find that the intensity of the spectral features (PAHs, emission lines and $9.7\mu\text{m}$ silicate absorption) of the extended components are very similar in all FEE_λ types, which suggests that the global physical properties of the star formation in the external parts (disks) of galaxies ($d \gtrsim 1.5\text{kpc}$) are likely the same. Moreover, the averaged spectra of the extended emission of all FEE_λ types resemble that of the template of lower IR luminosity starburst galaxies from Brandl et al. (2006). On the other hand, the spectra of the unresolved component are substantially different among the three




Figure 6. Weighted averaged spectra of the different FEE_λ types scaled to $13.2\mu\text{m}$ (unresolved: gray; constant/featureless: blue; PAH-dominated: green; silicate-dominated: red). The nuclear and extended emission components of each type are presented in the top and bottom panels respectively. On both panels we over-plot the local spectral template of Brandl et al. (2006) of normal star-forming galaxies in yellow, the averaged spectrum of the high-redshift submillimeter galaxies from Menéndez-Delmestre et al. (2009) in pink, as well as the template of the $z \sim 2$ ULIRGs from Farrah et al. (2008) in cyan. Unlike our GOALS galaxies, for which we show their nuclear and extended components, the comparison spectra are from the integrated emission of galaxies.

FEE_λ types even though the $f_{13.2\mu\text{m}}/f_{5.5\mu\text{m}}$ continuum ratio is similar in all of them, ranging between $\sim 5 - 6$. The nuclear MIR spectra of the GOALS galaxy sample will be analyzed in detail in a forthcoming work of Stierwalt et al. (2011, in prep.). From another point of view, these results also imply that, on average, the differences seen in the global/integrated MIR spectra of local ULIRGs and also LIRGs arise only from the processes that are taking place in their cores, while the extended star formation appear not to be affected by the nuclear activity and is similar to the more quiescent mode of star formation found in lower IR luminosity systems.

The difference between the MIR spectra of the unresolved emission of the constant and silicate-dominated FEE_λ types can be explained in terms of dust obscuration, with the silicate-dominated FEE_λ type having a larger $9.7\mu\text{m}$ optical depth. The relative emission of the PAHs with respect to that of the continuum in the unresolved spectrum of the PAH-dominated FEE_λ type is lower than in the constant/featureless FEE_λ type. This implies that a mechanism, such as an AGN or a compact nuclear starburst, is either destroying the PAH carriers in the nuclei of these galaxies, or probably simply diluting their strength (peak to continuum ratio) by heating the small dust grains to higher temperatures thus increasing the MIR continuum emission; a result that is in agreement with Sections 3.3 and 3.4.

In Figure 7 we show the averaged spectra of the total (blue), unresolved nuclear (red), and extended (green) emission for the galaxies in our sample corresponding

Table 1
Silicate Strengths and PAH EWs of the FEE $_{\lambda}$ Types

FEE $_{\lambda}$ Type (1)	Total		Unresolved		Extended	
	$S_{\text{Si } 9.7 \mu\text{m}}$ (2)	6.2 μm PAH EW (3)	$S_{\text{Si } 9.7 \mu\text{m}}$ (2)	6.2 μm PAH EW (3)	$S_{\text{Si } 9.7 \mu\text{m}}$ (2)	6.2 μm PAH EW (3)
Unresolved	-0.55 $\pm$ 0.01	0.05 $\pm$ 0.01	-0.31 $\pm$ 0.01	0.00 $\pm$ 0.01	...	...
Constant/featureless	-0.89 $\pm$ 0.01	0.50 $\pm$ 0.01	-1.01 $\pm$ 0.01	0.50 $\pm$ 0.02	-0.81 $\pm$ 0.02	0.54 $\pm$ 0.01
PAH-dominated	-0.74 $\pm$ 0.01	0.29 $\pm$ 0.01	-0.60 $\pm$ 0.03	0.24 $\pm$ 0.01	-0.77 $\pm$ 0.01	0.48 $\pm$ 0.01
Silicate-dominated	-1.53 $\pm$ 0.02	0.34 $\pm$ 0.02	-1.72 $\pm$ 0.02	0.43 $\pm$ 0.02	-0.94 $\pm$ 0.01	0.51 $\pm$ 0.01

Note. — (1) FEE $_{\lambda}$ type; (2) Silicate strength; (3) 6.2 μm PAH EW.




Figure 7. Averaged spectra of the total (blue), unresolved (red), and extended (green) emission components of our sample, normalized at 13.2 μm , for the different types of FEE $_{\lambda}$. The integrated spectral templates used to compare our sample with low and high-redshift star-forming galaxies are as in Figure 6. A total of 194 sources, for which both their FEE $_{13.2 \mu\text{m}}$ and $1 - \text{FEE}_{13.2 \mu\text{m}}$ are larger than 0.05 and therefore could be normalized at 13.2 μm , were used for this plot.

to each FEE $_{\lambda}$ type. In Table 1 we present the silicate strength ($S_{\text{Si } 9.7 \mu\text{m}}$) and 6.2 μm PAH EW of the different averaged spectra. The $S_{\text{Si } 9.7 \mu\text{m}}$ was calculated by fitting a power-law to the spectra, with anchors at 5.5, 6.6, and 13.2 μm , and evaluated at its maximum. The 6.2 μm PAH EW was obtained by fitting the local continuum of the feature using a linear function with anchors at 5.8 and 6.6 μm and integrating its emission from 5.9 to 6.5 μm .

By definition, the averaged spectra of the total as well as the unresolved and extended components for the con-

stant/featureless FEE $_{\lambda}$ type galaxies are very similar. The silicate strengths vary between -0.8 and -1.00 (see Table 1), and the 6.2 μm PAH EWs $\sim 0.5 \mu\text{m}$, which are close to the values found for the starburst galaxy sample of Brandl et al. (2006). The average spectrum of the extended emission component for the PAH-dominated FEE $_{\lambda}$ type, on the other hand, is very different from those of the unresolved and total components. Although the $S_{\text{Si } 9.7 \mu\text{m}}$ is similar, the 6.2 μm PAH EW of the extended spectrum (0.48 μm) is around twice that of the unresolved and total spectra (0.24 – 0.29 μm ; see Table 1). The average $f_{13.2 \mu\text{m}}/f_{5.5 \mu\text{m}}$ continuum ratios of the nuclear and extended spectra are also very similar ~ 7 . The averaged spectra of the components of the silicate-dominated FEE $_{\lambda}$ type are intrinsically different. The spectrum of the unresolved emission shows a deep silicate absorption feature ($S_{\text{Si } 9.7 \mu\text{m}} = -1.72$) while the spectrum of the extended emission is significantly shallower ($S_{\text{Si } 9.7 \mu\text{m}} = -0.94$) but similar to the values found for the extended components of the constant/featureless and PAH-dominated FEE $_{\lambda}$ types. If we define $\Delta S_{\text{Si } 9.7 \mu\text{m}}$ as the difference between the silicate strength of the nuclear and extended component of a galaxy, then $\Delta S_{\text{Si } 9.7 \mu\text{m}} = -0.78$ for the silicate-dominated FEE $_{\lambda}$ type (see also Section 3.2). The averaged nuclear spectra of the unresolved FEE $_{\lambda}$ type (Figure 6) show an almost flat rising continuum, a moderate silicate strength ($S_{\text{Si } 9.7 \mu\text{m}} = -0.31$), and almost no PAH emission (6.2 μm PAH EW $\sim 0.00 \mu\text{m}$), all of them typical signatures of AGN-dominated galaxies. Their MIR spectrum is similar to sources in region 1A of Spoon et al. (2007) diagram, in which the AGN emission is the likely culprit for the dilution of the PAH features.

The classification of the GOALS galaxies into different stages of interaction carried out by Petric et al. (2011) and Haan et al. (2011) also allows us to explore whether the PAH-to-continuum FEE ratios or the MIR spectral shape of the nuclear and extended components of galaxies are related to the merger stage of the systems. Unfortunately, no such clear trend is found. Only at the final stage of interaction (stage 4), when the nuclei of the galaxies are already assembled, the averaged spectrum of their unresolved component shows a deeper silicate absorption feature than that displayed by the spectrum of the extended component. This is understood if we take into account that most of the ULIRGs in GOALS, which show the largest silicate strengths, are classified as being in this latest merger stage. On the other hand, the averaged spectra of both components for galaxies grouped in stages 0 to 3 are very similar, and resemble that of local starburst galaxies.

3.6. High-redshift Implications

In Figures 6 and 7 we also compare the averaged spectra of the GOALS sample, both nuclear and extended components, with the composite, integrated spectra of the $z \sim 2$ ULIRG sample from Farrah et al. (2008), and the sub-millimeter galaxies (SMGs) studied in Menéndez-Delmestre et al. (2009) which are located at a redshift range of $z \sim 0.65 - 3.2$. As already mentioned in Farrah et al. (2008), their composite spectrum of high- z ULIRGs and HyperLIRGs ($L_{\text{IR}} = 10^{12.9} - 10^{13.8} L_{\odot}$) is very similar to that of the starburst galaxies from Brandl et al. (2006). We also note that it is also very similar to the averaged spectra of the extended components of our galaxy sample. This suggests that the properties of star forming high- z ULIRGs, often selected based on the characteristic *Spitzer*/IRAC colors due to the $1.6 \mu\text{m}$ stellar bump and their *Spitzer*/MIPS $24 \mu\text{m}$ fluxes (e.g., Magdis et al. 2010) is very similar to the *extended disks* of local (U)LIRGs. However, the composite, integrated SMG spectrum is different from our average spectrum of the extended component, displaying PAHs with a considerably lower EW. Menéndez-Delmestre et al. (2009) argued that an additional power-law-type source of hot dust emission was needed to partially enhance the MIR continuum of these galaxies. They suggested that this would arise either from an AGN torus or from optically thick dust emission around star-forming regions, in agreement with the scenario we propose for the compact nuclear sources in the previous Sections. Interestingly, the SMG composite spectrum is more similar to the averaged nuclear spectrum of (U)LIRGs belonging to the PAH-dominated FEE $_{\lambda}$ galaxy type (see Stierwalt et al. 2011, in prep.). We also note that while the [Ne II] $12.81 \mu\text{m}$ emission line is clearly visible in all of our averaged spectra as well as in the high- z ULIRG spectrum, it is almost absent from that of the SMGs. Nevertheless, we attribute this difference to the limited quality of the composite spectrum of the SMGs, which prevent us from performing meaningful comparisons regarding specific features in the spectra.

Summarizing, the spectra of the extended components of (U)LIRGs are very similar independently of their FEE $_{\lambda}$ type, suggesting that the extended disk emission of all LIRGs and ULIRGs could be dominated by PAHs excited principally by non-ionizing stars and quiescent star-forming regions. The fact that recent *Herschel* observations imply that the galactic disks are responsible for the bulk of the diffuse cold dust emission peaking at $100 - 500 \mu\text{m}$ (Bendo et al. 2010) leads us to speculate whether this would also hold in (U)LIRGs. Within this picture, their outer disks, traced by the PAH spatial profiles, would be responsible for the cold FIR emission. Meanwhile, the inner nuclei of the most compact sources, where most of their energy is produced, would dominate the MIR continuum and also contribute to show warmer FIR colors than galaxies without compact cores. Due to their distance and small physical size this can not be verified directly by *Herschel* or even future missions, such as SPICA. However, modeling the FIR line emission (e.g., van der Werf et al. 2010; González-Alfonso et al. 2010) will enable us to probe the excitation and density of gas and thus provide further constraints on its spatial distribution.

4. CONCLUSIONS

We analyzed the spatial profiles of low spectral resolution $5 - 14 \mu\text{m}$ *Spitzer*/IRS spectra of the GOALS galaxy sample and quantified the spatial extent of their MIR emission, FEE $_{\lambda}$. Our work indicates that:

- The [Ne II] $12.81 \mu\text{m}$ emission is as compact as the MIR continuum, but the PAHs are more extended in many galaxies. This is in agreement with studies showing that the [Ne II] $12.81 \mu\text{m}$ and MIR continuum emissions trace the ionizing stars (i.e., clumpy, current star formation), while the PAH emission is more representative of older, colder, and diffuse stellar populations.
- The $11.3 \mu\text{m}$ PAH emission is more extended than that of the 6.2 and $7.7 \mu\text{m}$ PAHs, which is consistent with the formers being enhanced in a more ionized medium while the $11.3 \mu\text{m}$ PAH emission is similar for neutral and ionized molecules.
- On average, the MIR continuum becomes more compact than the PAH emission as the MIR is increasingly dominated by an AGN. That is, the presence of an AGN modifies the spatial extent of the hot dust continuum but has a negligible effect on the overall, kpc-scale distribution of the PAH emission. None of the AGN-dominated (U)LIRGs shows a constant/featureless FEE $_{\lambda}$ type, implying that the AGN leaves a “footprint” in the FEE $_{\lambda}$ function of a galaxy. Therefore, the MIR emission of the galaxies showing a constant FEE $_{\lambda}$ will not be dominated by an AGN or, in general, show compact MIR continuum emission.
- The PAH-to-continuum FEE ratio of (U)LIRGs increases as the global dust temperature, traced by their *IRAS* $\log(f_{60 \mu\text{m}}/f_{100 \mu\text{m}})$ FIR color, becomes warmer. This can be attributed to the presence of a compact, powerful (circum-)nuclear starburst in these galaxies which would increase the temperature of the dust, and also the ratio between the volume of the shells of the photo-dissociation regions –from where PAH emission arises– and the volume of the hot dust emitting region. “Normal” galaxies do not show enhanced PAH-to-continuum FEE ratios but a rather constant value. On the other hand, in (U)LIRGs with large PAH-to-continuum FEE ratios the MIR continuum is more compact than the PAH emission, which is still as extended as in the “normal” population since it arises mostly from a diffuse component (their disks). Nonetheless, in terms of the relative spatial distribution of the PAH and continuum emissions, this effect is indistinguishable from that caused by an AGN. Moreover, both processes, *compact* star formation and AGN activity, coexist in 27% of our (U)LIRGs sample.
- The intensity of the spectral features and continuum emission of the extended emission component of all (U)LIRGs are very similar, indicating that the properties of the star formation in the external parts of galaxies (disks; $d \gtrsim 1.5 \text{ kpc}$) are uniform.

Moreover, they are similar to local starburst galaxies as well as to $z \sim 2$ ULIRGs (selected by their *Spitzer*/IRAC and MIPS colors), but different from sub-millimeter-selected high-redshift galaxies. On the other hand, the MIR spectra of the unresolved nuclear component of galaxies vary widely.

- These results imply that, on average, the diversity seen in the properties of global/integrated MIR spectra of local ULIRGs as well as LIRGs arise only from the processes that are taking place in their cores. The extended star formation appears unaffected by the nuclear activity and it is similar to the more quiescent mode of star formation found in lower IR luminosity systems. We speculate that the outer disks/regions of (U)LIRGs, traced by their PAH spatial profiles, are responsible for most of the cold FIR emission as seen in galaxies with lower IR luminosities, while the nuclei of the most compact (U)LIRGs, as stated in Díaz-Santos et al. (2010b), would dominate the MIR continuum and contribute to show overall warmer FIR colors.

We thank the referee for her/his useful comments that helped to improve the paper. TDS would like to thank D. Elbaz, E. Le Floc'h, V. Leboutellier, E. Daddi, and G. Magdis for stimulating discussions, as well as all colleagues at CEA/Saclay (France), where part of this work was done, for their hospitality. TDS and VC would like to acknowledge partial support from the EU ToK grant 39965 and FP7-REGPOT 206469. This research has made use of the NASA/IPAC Extragalactic Database (NED), which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration, and of NASA's Astrophysics Data System (ADS) abstract service.

REFERENCES

- Allamandola, L. J., Hudgins, D. M., & Sandford, S. A. 1999, *ApJ*, 511, L115
- Alonso-Herrero, A., Rieke, G. H., Rieke, M. J., & Scoville, N. Z. 2002, *AJ*, 124, 166
- Armus, L., et al. 2007, *ApJ*, 656, 148
- . 2009, *PASP*, 121, 559
- Bendo, G. J., et al. 2010, *A&A*, 518, L65+
- Brandl, B. R., et al. 2006, *ApJ*, 653, 1129
- Calzetti, D., et al. 2007, *ApJ*, 666, 870
- Caputi, K. I., et al. 2007, *ApJ*, 660, 97
- Charmandaris, V., Le Floc'h, E., & Mirabel, I. F. 2004, *ApJ*, 600, L15
- Charmandaris, V., et al. 2002, *A&A*, 391, 429
- Díaz-Santos, T., Alonso-Herrero, A., Colina, L., Packham, C., Levenson, N. A., Pereira-Santaella, M., Roche, P. F., & Telesco, C. M. 2010a, *ApJ*, 711, 328
- Díaz-Santos, T., Alonso-Herrero, A., Colina, L., Packham, C., Radomski, J. T., & Telesco, C. M. 2008, *ApJ*, 685, 211
- Díaz-Santos, T., et al. 2010b, *ApJ*, 723, 993
- Draine, B. T., & Li, A. 2001, *ApJ*, 551, 807
- Farrah, D., et al. 2008, *ApJ*, 677, 957
- Galliano, F., Madden, S. C., Tielens, A. G. G. M., Peeters, E., & Jones, A. P. 2008, *ApJ*, 679, 310
- García-Marín, M., Colina, L., & Arribas, S. 2009, *A&A*, 505, 1017
- Genzel, R., et al. 1998, *ApJ*, 498, 579
- González-Alfonso, E., et al. 2010, *A&A*, 518, L43+
- Haan, S., et al. 2011, *AJ*, 141, 100
- Howell, J. H., et al. 2010, *ApJ*, 715, 572
- Imanishi, M. 2009, *ApJ*, 694, 751
- Imanishi, M., Dudley, C. C., Maiolino, R., Maloney, P. R., Nakagawa, T., & Risaliti, G. 2007, *ApJS*, 171, 72
- Imanishi, M., Maiolino, R., & Nakagawa, T. 2010, *ApJ*, 709, 801
- Kramer, C., et al. 2010, *A&A*, 518, L67+
- Laurent, O., Mirabel, I. F., Charmandaris, V., Gallais, P., Madden, S. C., Sauvage, M., Vigroux, L., & Cesarsky, C. 2000, *A&A*, 359, 887
- Le Floc'h, E., et al. 2005, *ApJ*, 632, 169
- Lutz, D., Spoon, H. W. W., Rigopoulou, D., Moorwood, A. F. M., & Genzel, R. 1998b, *ApJ*, 505, L103
- Magdis, G. E., et al. 2010, *MNRAS*, 409, 22
- Magnelli, B., Elbaz, D., Chary, R. R., Dickinson, M., Le Borgne, D., Frayer, D. T., & Willmer, C. N. A. 2011, *A&A*, 528, A35+
- Menéndez-Delmestre, K., et al. 2009, *ApJ*, 699, 667
- Murphy, E. J., Chary, R., Dickinson, M., Pope, A., Frayer, D. T., & Lin, L. 2011, *ArXiv e-prints*
- Netzer, H., et al. 2007, *ApJ*, 666, 806
- Pereira-Santaella, M., Alonso-Herrero, A., Rieke, G. H., Colina, L., Díaz-Santos, T., & et al. 2010, *ApJ*, in press
- Pérez-González, P. G., et al. 2005, *ApJ*, 630, 82
- Petric, A. O., et al. 2011, *ApJ*, 730, 28
- Sales, D. A., Pastoriza, M. G., & Riffel, R. 2010, *ApJ*, 725, 605
- Sanders, D. B., Mazzarella, J. M., Kim, D.-C., Surace, J. A., & Soifer, B. T. 2003, *AJ*, 126, 1607
- Sanders, D. B., & Mirabel, I. F. 1996, *ARA&A*, 34, 749
- Sauvage, M., & Thuan, T. X. 1992, *ApJ*, 396, L69
- Sauvage, M., et al. 2010, *A&A*, 518, L64+
- Schweitzer, M., et al. 2006, *ApJ*, 649, 79
- Smith, J. D. T., et al. 2004, *ApJS*, 154, 199
- . 2007, *ApJ*, 656, 770
- Soifer, B. T., Bock, J. J., Marsh, K., Neugebauer, G., Matthews, K., Egami, E., & Armus, L. 2003, *AJ*, 126, 143
- Soifer, B. T., Neugebauer, G., Matthews, K., Egami, E., & Weinberger, A. J. 2002, *AJ*, 124, 2980
- Spoon, H. W. W., Marshall, J. A., Houck, J. R., Elitzur, M., Hao, L., Armus, L., Brandl, B. R., & Charmandaris, V. 2007, *ApJ*, 654, L49
- van der Werf, P. P., et al. 2010, *A&A*, 518, L42+
- Veilleux, S., Kim, D., & Sanders, D. B. 2002, *ApJS*, 143, 315
- Veilleux, S., et al. 2009, *ApJS*, 182, 628
- Walterbos, R. A. M., & Greenawalt, B. 1996, *ApJ*, 460, 696
- Wu, Y., Charmandaris, V., Huang, J., Spinoglio, L., & Tommasin, S. 2009, *ApJ*, 701, 658

Table 2
FEEs of Various MIR Features

Galaxy name (1)	R.A. (J2000) (2)	Declination (J2000) (3)	FEE _{6.2μm} (PAH) (4)	FEE _{6.7μm} (Cont.) (5)	FEE _{7.7μm} (PAH) (6)	FEE _{9.7μm} (PAH) (7)	FEE _{11.3μm} (Si abs.) (8)	FEE _{12.8μm} ([Ne II]) (9)	FEE $_{\lambda}$ Type (10)
NGC0023	00h 09m 53.35s	+25° 55m 27.8s	0.66 ± 0.02	0.65 ± 0.01	0.68 ± 0.02	0.66 ± 0.01	0.67 ± 0.03	0.68 ± 0.02	1
MCG-02-01-051	00h 18m 50.90s	−10° 22m 36.8s	0.37 ± 0.04	0.38 ± 0.02	0.36 ± 0.05	0.40 ± 0.02	0.29 ± 0.06	0.43 ± 0.03	1
ESO350-IG038	00h 36m 52.49s	−33° 33m 17.3s	0.08 ± 0.04	0.12 ± 0.02	0.13 ± 0.04	0.17 ± 0.04	0.10 ± 0.06	0.17 ± 0.04	2
NGC0232	00h 42m 45.84s	−23° 33m 41.0s	0.36 ± 0.07	0.40 ± 0.01	0.49 ± 0.04	0.41 ± 0.01	0.38 ± 0.05	0.47 ± 0.04	3
NGC0232	00h 42m 52.82s	−23° 32m 28.0s	0.28 ± 0.09	0.35 ± 0.02	0.29 ± 0.07	0.35 ± 0.02	0.22 ± 0.06	0.37 ± 0.04	2
MCG+12-02-001	00h 54m 03.89s	+73° 05m 06.0s	0.47 ± 0.02	0.47 ± 0.04	0.32 ± 0.04	0.50 ± 0.01	0.44 ± 0.05	0.49 ± 0.04	2
NGC0317B	00h 57m 40.42s	+43° 47m 32.6s	0.25 ± 0.06	0.24 ± 0.01	0.36 ± 0.06	0.29 ± 0.07	0.19 ± 0.07	0.33 ± 0.04	3
MCG-03-04-014	01h 10m 08.93s	−16° 51m 10.1s	0.50 ± 0.04	0.53 ± 0.01	0.58 ± 0.03	0.54 ± 0.01	0.53 ± 0.04	0.57 ± 0.03	1
ESO244-G012	01h 18m 08.23s	−44° 28m 00.5s	0.16 ± 0.05	0.18 ± 0.02	0.20 ± 0.03	0.20 ± 0.01	0.12 ± 0.05	0.20 ± 0.04	1
CGCG436-030	01h 20m 02.64s	+14° 21m 42.1s	0.17 ± 0.06	0.26 ± 0.03	0.36 ± 0.06	0.24 ± 0.05	0.07 ± 0.07	0.35 ± 0.05	3
ESO353-G020	01h 34m 51.26s	−36° 08m 14.3s	0.36 ± 0.04	0.41 ± 0.02	0.53 ± 0.04	0.44 ± 0.01	0.44 ± 0.05	0.50 ± 0.04	3
ESO297-G011	01h 36m 23.40s	−37° 19m 18.1s	0.52 ± 0.04	0.53 ± 0.02	0.51 ± 0.04	0.56 ± 0.01	0.53 ± 0.05	0.58 ± 0.04	1
ESO297-G011	01h 36m 24.14s	−37° 20m 25.8s	0.16 ± 0.07	0.16 ± 0.01	0.21 ± 0.07	0.19 ± 0.03	0.09 ± 0.06	0.21 ± 0.05	1
IRASF01364-1042	01h 38m 52.80s	−10° 27m 12.2s	0.09 ± 0.10	0.08 ± 0.04	0.24 ± 0.06	0.16 ± 0.10	0.06 ± 0.08	0.23 ± 0.07	2
IRASF01417+1651	01h 44m 30.55s	+17° 06m 09.0s	0.35 ± 0.06	0.28 ± 0.09	0.57 ± 0.13	0.42 ± 0.05	0.41 ± 0.07	0.49 ± 0.03	3
NGC0695	01h 51m 14.35s	+22° 34m 55.9s	0.71 ± 0.01	0.74 ± 0.01	0.74 ± 0.02	0.74 ± 0.01	0.73 ± 0.03	0.75 ± 0.02	1
UGC01385	01h 54m 53.83s	+36° 55m 04.4s	0.29 ± 0.07	0.25 ± 0.07	0.29 ± 0.03	0.28 ± 0.07	0.23 ± 0.05	0.29 ± 0.04	1
NGC0838	02h 09m 42.82s	−10° 11m 02.0s	0.11 ± 0.05	0.19 ± 0.01	0.30 ± 0.05	0.19 ± 0.02	0.11 ± 0.06	0.31 ± 0.04	2
NGC0838	02h 09m 38.66s	−10° 08m 47.0s	0.54 ± 0.04	0.54 ± 0.01	0.59 ± 0.03	0.55 ± 0.01	0.58 ± 0.04	0.61 ± 0.03	1
NGC0838	02h 09m 20.88s	−10° 07m 59.5s	0.73 ± 0.01	0.75 ± 0.02	0.72 ± 0.02	0.74 ± 0.01	0.74 ± 0.03	0.74 ± 0.02	1
NGC0828	02h 10m 09.53s	+39° 11m 24.7s	0.70 ± 0.01	0.71 ± 0.02	0.74 ± 0.02	0.72 ± 0.01	0.72 ± 0.03	0.73 ± 0.02	1
IC0214	02h 14m 05.57s	+05° 10m 23.9s	0.72 ± 0.02	0.72 ± 0.01	0.72 ± 0.02	0.72 ± 0.01	0.72 ± 0.03	0.73 ± 0.03	1
NGC0877	02h 17m 59.69s	+14° 32m 38.0s	0.80 ± 0.05	0.79 ± 0.06	0.79 ± 0.07	0.76 ± 0.04	0.80 ± 0.06	0.80 ± 0.06	1
NGC0877	02h 17m 53.26s	+14° 31m 18.5s	0.41 ± 0.04	0.39 ± 0.03	0.58 ± 0.04	0.34 ± 0.05	0.33 ± 0.05	0.50 ± 0.03	3
MCG+05-06-036	02h 23m 21.98s	+32° 11m 48.8s	0.29 ± 0.05	0.33 ± 0.03	0.35 ± 0.04	0.36 ± 0.05	0.26 ± 0.06	0.35 ± 0.04	1
MCG+05-06-036	02h 23m 18.96s	+32° 11m 18.6s	0.50 ± 0.03	0.47 ± 0.02	0.54 ± 0.03	0.51 ± 0.01	0.49 ± 0.05	0.55 ± 0.03	1
UGC01845	02h 24m 07.97s	+47° 58m 12.0s	0.49 ± 0.03	0.50 ± 0.04	0.56 ± 0.02	0.53 ± 0.01	0.53 ± 0.04	0.56 ± 0.03	1
NGC0992	02h 37m 25.46s	+21° 06m 02.9s	0.75 ± 0.02	0.74 ± 0.01	0.76 ± 0.02	0.74 ± 0.01	0.76 ± 0.02	0.76 ± 0.02	1
UGC02238	02h 46m 17.45s	+13° 05m 44.5s	0.67 ± 0.02	0.67 ± 0.01	0.74 ± 0.02	0.67 ± 0.01	0.67 ± 0.03	0.72 ± 0.02	3
IRASF02437+2122	02h 46m 39.12s	+21° 35m 10.3s	0.14 ± 0.05	0.11 ± 0.02	0.37 ± 0.11	0.11 ± 0.02	0.04 ± 0.06	0.15 ± 0.07	3
UGC02369	02h 54m 01.75s	+14° 58m 36.5s	0.49 ± 0.04	0.52 ± 0.02	0.37 ± 0.04	0.53 ± 0.01	0.50 ± 0.04	0.53 ± 0.03	2
UGC02608	03h 15m 01.46s	+42° 02m 08.5s	0.17 ± 0.03	0.21 ± 0.05	0.26 ± 0.03	0.28 ± 0.05	0.22 ± 0.05	0.29 ± 0.04	2
UGC02608	03h 15m 14.57s	+41° 58m 50.2s	0.41 ± 0.01	0.55 ± 0.18	0.47 ± 0.01	0.57 ± 0.07	0.70 ± 0.06	0.72 ± 0.04	2
NGC1275	03h 19m 48.17s	+41° 30m 42.1s	0.09 ± 0.07	0.14 ± 0.01	0.04 ± 0.04	0.14 ± 0.06	0.01 ± 0.01	0.02 ± 0.06	0
IRASF03217+4022	03h 25m 05.38s	+40° 33m 32.0s	0.46 ± 0.02	0.49 ± 0.03	0.52 ± 0.03	0.51 ± 0.01	0.42 ± 0.05	0.53 ± 0.04	1
NGC1365	03h 33m 36.41s	−36° 08m 25.8s	0.47 ± 0.01	0.70 ± 0.03	0.30 ± 0.10	0.74 ± 0.01	0.65 ± 0.03	0.67 ± 0.02	2
IRASF03359+1523	03h 38m 47.06s	+15° 32m 54.2s	... ± ...	0.33 ± 0.11	0.41 ± 0.09	0.38 ± 0.02	0.63 ± 0.04	0.59 ± 0.03	4
CGCG465-012	03h 54m 07.68s	+15° 59m 24.4s	0.72 ± 0.03	0.73 ± 0.01	0.72 ± 0.05	0.71 ± 0.02	0.72 ± 0.04	0.72 ± 0.04	1
CGCG465-012	03h 54m 15.96s	+15° 55m 43.3s	0.58 ± 0.03	0.59 ± 0.01	0.61 ± 0.03	0.59 ± 0.01	0.57 ± 0.04	0.64 ± 0.03	1
IRAS03582+6012	04h 02m 33.00s	+60° 20m 41.6s	0.02 ± 0.01	0.10 ± 0.08	0.49 ± 0.05	0.07 ± 0.04	0.12 ± 0.07	0.42 ± 0.04	3
IRAS03582+6012	04h 02m 31.97s	+60° 20m 38.4s	0.50 ± 0.04	0.49 ± 0.01	0.53 ± 0.03	0.49 ± 0.01	0.47 ± 0.05	0.52 ± 0.04	1
UGC02982	04h 12m 22.68s	+05° 32m 49.2s	0.66 ± 0.01	0.66 ± 0.02	0.72 ± 0.01	0.67 ± 0.02	0.69 ± 0.03	0.72 ± 0.02	1
ESO420-G013	04h 13m 49.70s	−32° 00m 25.2s	0.25 ± 0.04	0.32 ± 0.05	0.23 ± 0.07	0.38 ± 0.01	0.29 ± 0.05	0.38 ± 0.04	2
NGC1572	04h 22m 42.82s	−40° 36m 03.2s	0.31 ± 0.05	0.38 ± 0.01	0.44 ± 0.03	0.39 ± 0.01	0.39 ± 0.05	0.43 ± 0.04	1
IRAS04271+3849	04h 30m 33.10s	+38° 55m 47.6s	0.32 ± 0.05	0.37 ± 0.01	0.46 ± 0.04	0.39 ± 0.01	0.35 ± 0.05	0.42 ± 0.05	3
NGC1614	04h 33m 59.95s	−08° 34m 46.6s	0.33 ± 0.07	0.34 ± 0.10	0.40 ± 0.04	0.43 ± 0.01	0.36 ± 0.05	0.45 ± 0.04	1
UGC03094	04h 35m 33.82s	+19° 10m 18.1s	0.73 ± 0.01	0.78 ± 0.02	0.72 ± 0.02	0.78 ± 0.01	0.76 ± 0.02	0.78 ± 0.02	2

Note. — (1) Galaxy name. Multiple systems are indicated with the same name but providing the right ascension and declination of the individual galaxies. An asterisk next to a name indicates that multiple nuclei, not resolved by *Spitzer*/IRS, are detected using higher angular resolution *HST* NIR continuum imaging (Haan et al. 2011). (2) Right Ascension (J2000). (3) Declination (J2000). (4)–(9) Fraction of extended emission (FEE) at: 6.2, 6.7, 7.7, 9.7, 11.3, and 12.8 μ m respectively. (10) FEE $_{\lambda}$ function type: 0, FEE $_{\lambda} \sim 0$; 1, constant/featureless; 2, PAH-dominated; 3, silicate-dominated; 4, undefined.

Table 2
FEEs of Various MIR Features

Galaxy name (1)	R.A. (J2000) (2)	Declination (J2000) (3)	FEE _{6.2μm} (PAH) (4)	FEE _{6.7μm} (Cont.) (5)	FEE _{7.7μm} (PAH) (6)	FEE _{9.7μm} (PAH) (7)	FEE _{11.3μm} (Si abs.) (8)	FEE _{12.8μm} ([Ne II]) (9)	FEE $_{\lambda}$ Type (10)
ESO203-IG001*	04h 46m 49.56s	-48° 33m 30.6s	0.12 ± 0.07	0.16 ± 0.02	0.39 ± 0.14	0.22 ± 0.04	0.05 ± 0.01	0.30 ± 0.07	3
MCG-05-12-006	04h 52m 04.97s	-32° 59m 26.2s	0.08 ± 0.08	0.20 ± 0.01	0.18 ± 0.05	0.20 ± 0.01	0.10 ± 0.06	0.21 ± 0.05	1
NGC1797	05h 07m 44.83s	-08° 01m 08.8s	0.25 ± 0.06	0.29 ± 0.02	0.30 ± 0.04	0.26 ± 0.08	0.29 ± 0.05	0.33 ± 0.03	1
CGCG468-002	05h 08m 19.70s	+17° 21m 47.9s	0.14 ± 0.09	0.24 ± 0.02	0.10 ± 0.06	0.26 ± 0.03	0.14 ± 0.06	0.20 ± 0.04	2
CGCG468-002	05h 08m 21.22s	+17° 22m 08.0s	0.39 ± 0.06	0.40 ± 0.01	0.52 ± 0.04	0.41 ± 0.02	0.28 ± 0.06	0.49 ± 0.04	3
IRAS05083+2441	05h 11m 25.87s	+24° 45m 18.4s	0.39 ± 0.06	0.39 ± 0.03	0.45 ± 0.04	0.42 ± 0.01	0.37 ± 0.05	0.44 ± 0.04	1
IRASF05081+7936	05h 16m 46.39s	+79° 40m 13.1s	0.42 ± 0.04	0.43 ± 0.02	0.49 ± 0.04	0.45 ± 0.01	0.41 ± 0.07	0.48 ± 0.04	1
IRAS05129+5128	05h 16m 55.97s	+51° 31m 57.0s	0.18 ± 0.07	0.34 ± 0.01	0.39 ± 0.06	0.34 ± 0.03	0.25 ± 0.06	0.37 ± 0.05	1
IRASF05189-2524	05h 21m 01.44s	-25° 21m 46.1s	... ± ...	0.17 ± 0.04	0.07 ± 0.01	0.18 ± 0.04	... ± ...	... ± ...	0
IRASF05187-1017	05h 21m 06.53s	-10° 14m 46.3s	0.15 ± 0.03	0.20 ± 0.07	0.47 ± 0.12	0.25 ± 0.05	0.15 ± 0.07	0.31 ± 0.06	3
IRAS05223+1908	05h 25m 16.68s	+19° 10m 47.6s	0.05 ± 0.06	0.04 ± 0.01	0.09 ± 0.09	0.10 ± 0.05	0.02 ± 0.01	0.03 ± 0.01	0
MCG+08-11-002	05h 40m 43.70s	+49° 41m 41.6s	0.26 ± 0.05	0.27 ± 0.01	0.56 ± 0.03	0.27 ± 0.08	0.28 ± 0.06	0.47 ± 0.03	3
NGC1961	05h 42m 04.56s	+69° 22m 43.0s	0.66 ± 0.05	0.70 ± 0.03	0.71 ± 0.02	0.72 ± 0.01	0.71 ± 0.02	0.74 ± 0.02	1
UGC03351	05h 45m 48.02s	+58° 42m 03.6s	0.74 ± 0.01	0.78 ± 0.02	0.81 ± 0.02	0.78 ± 0.02	0.75 ± 0.03	0.79 ± 0.02	1
IRAS05442+1732	05h 47m 11.21s	+17° 33m 46.4s	0.28 ± 0.03	0.24 ± 0.07	0.29 ± 0.03	0.29 ± 0.06	0.25 ± 0.06	0.34 ± 0.04	1
IRASF06076-2139	06h 09m 45.74s	-21° 40m 24.6s	0.16 ± 0.05	0.17 ± 0.02	0.27 ± 0.05	0.22 ± 0.05	0.09 ± 0.10	0.22 ± 0.05	3
UGC03410	06h 14m 29.62s	+80° 26m 59.6s	0.75 ± 0.01	0.75 ± 0.01	0.77 ± 0.02	0.75 ± 0.01	0.76 ± 0.03	0.77 ± 0.02	1
UGC03410	06h 13m 57.89s	+80° 28m 34.7s	0.68 ± 0.01	0.67 ± 0.01	0.73 ± 0.02	0.72 ± 0.02	0.73 ± 0.02	0.74 ± 0.01	1
NGC2146	06h 18m 37.82s	+78° 21m 24.1s	0.60 ± 0.03	0.60 ± 0.01	0.70 ± 0.02	0.60 ± 0.02	0.65 ± 0.03	0.68 ± 0.02	3
ESO255-IG007	06h 27m 21.70s	-47° 10m 36.1s	0.17 ± 0.08	0.24 ± 0.01	0.24 ± 0.05	0.26 ± 0.01	0.16 ± 0.06	0.27 ± 0.04	1
ESO255-IG007	06h 27m 22.56s	-47° 10m 47.3s	0.54 ± 0.07	0.49 ± 0.02	0.48 ± 0.04	0.55 ± 0.01	0.54 ± 0.04	0.56 ± 0.04	1
ESO255-IG007	06h 27m 23.09s	-47° 11m 02.8s	0.27 ± 0.17	0.26 ± 0.03	0.25 ± 0.13	0.30 ± 0.01	0.24 ± 0.06	0.36 ± 0.04	1
ESO557-G002	06h 31m 47.21s	-17° 37m 16.7s	0.37 ± 0.21	0.26 ± 0.20	0.58 ± 0.03	0.48 ± 0.04	0.43 ± 0.05	0.56 ± 0.03	3
UGC03608	06h 57m 34.42s	+46° 24m 10.4s	0.57 ± 0.05	0.56 ± 0.04	0.43 ± 0.04	0.54 ± 0.06	0.37 ± 0.05	0.54 ± 0.02	2
IRASF06592-6313	06h 59m 40.25s	-63° 17m 52.4s	0.10 ± 0.04	0.17 ± 0.06	0.20 ± 0.02	0.17 ± 0.07	0.11 ± 0.05	0.18 ± 0.04	1
IRASF07027-6011*	07h 03m 28.51s	-60° 16m 43.7s	0.36 ± 0.04	0.34 ± 0.04	0.40 ± 0.04	0.40 ± 0.01	0.37 ± 0.06	0.43 ± 0.05	1
NGC2342	07h 09m 18.07s	+20° 38m 10.3s	0.31 ± 0.03	0.33 ± 0.02	0.39 ± 0.03	0.33 ± 0.04	0.29 ± 0.05	0.38 ± 0.04	1
IRAS07251-0248	07h 27m 37.63s	-02° 54m 54.7s	0.14 ± 0.05	0.10 ± 0.07	0.32 ± 0.10	0.13 ± 0.06	0.09 ± 0.01	0.14 ± 0.06	3
NGC2388	07h 29m 04.58s	+33° 51m 38.2s	0.33 ± 0.03	0.32 ± 0.03	0.32 ± 0.03	0.36 ± 0.02	0.30 ± 0.05	0.39 ± 0.04	1
MCG+02-20-003	07h 35m 43.44s	+11° 42m 34.9s	0.12 ± 0.06	0.30 ± 0.01	0.51 ± 0.02	0.28 ± 0.08	0.31 ± 0.05	0.47 ± 0.03	3
IRAS08355-4944	08h 37m 01.87s	-49° 54m 29.9s	0.10 ± 0.06	0.26 ± 0.01	0.18 ± 0.07	0.30 ± 0.09	0.24 ± 0.06	0.30 ± 0.03	2
NGC2623	08h 38m 24.12s	+25° 45m 16.6s	0.21 ± 0.03	0.22 ± 0.04	0.42 ± 0.03	0.26 ± 0.06	0.13 ± 0.08	0.30 ± 0.05	3
ESO432-IG006	08h 44m 27.22s	-31° 41m 50.6s	0.37 ± 0.06	0.42 ± 0.01	0.45 ± 0.04	0.42 ± 0.01	0.42 ± 0.05	0.45 ± 0.04	1
ESO432-IG006	08h 44m 28.92s	-31° 41m 30.1s	0.20 ± 0.06	0.20 ± 0.06	0.28 ± 0.05	0.16 ± 0.09	0.13 ± 0.06	0.19 ± 0.04	1
ESO60-IG016	08h 52m 30.50s	-69° 01m 59.2s	0.11 ± 0.01	0.24 ± 0.07	0.36 ± 0.05	0.22 ± 0.06	0.18 ± 0.06	0.33 ± 0.05	3
IRASF08572+3915	09h 00m 25.34s	+39° 03m 54.0s	0.04 ± 0.02	0.16 ± 0.06	0.17 ± 0.17	0.10 ± 0.09	... ± ...	0.06 ± 0.05	0
IRAS09022-3615	09h 04m 12.70s	-36° 27m 01.4s	0.10 ± 0.03	0.13 ± 0.03	0.17 ± 0.04	0.20 ± 0.02	0.10 ± 0.08	0.20 ± 0.05	2
IRASF09111-1007	09h 13m 36.50s	-10° 19m 29.6s	0.24 ± 0.08	0.28 ± 0.02	0.35 ± 0.04	0.27 ± 0.04	0.18 ± 0.06	0.30 ± 0.04	1
IRASF09111-1007	09h 13m 38.88s	-10° 19m 19.6s	0.43 ± 0.12	0.42 ± 0.04	0.39 ± 0.04	0.42 ± 0.01	0.32 ± 0.05	0.43 ± 0.04	1
UGC04881	09h 15m 55.51s	+44° 19m 57.4s	0.37 ± 0.03	0.52 ± 0.03	0.59 ± 0.05	0.49 ± 0.03	0.35 ± 0.06	0.57 ± 0.04	3
UGC04881	09h 15m 54.70s	+44° 19m 50.9s	0.33 ± 0.05	0.38 ± 0.02	0.48 ± 0.03	0.37 ± 0.03	0.40 ± 0.06	0.44 ± 0.04	3
UGC05101	09h 35m 51.60s	+61° 21m 11.9s	0.21 ± 0.07	0.30 ± 0.02	0.35 ± 0.03	0.28 ± 0.02	0.20 ± 0.05	0.36 ± 0.04	2
MCG+08-18-013	09h 36m 37.20s	+48° 28m 27.8s	... ± ...	0.42 ± 0.01	... ± ...	0.47 ± 0.07	0.50 ± 0.01	0.56 ± 0.11	4
IRASF09437+0317	09h 46m 21.10s	+03° 04m 16.3s	0.64 ± 0.02	0.60 ± 0.04	0.67 ± 0.07	0.62 ± 0.03	0.64 ± 0.06	0.66 ± 0.06	1
IRASF09437+0317	09h 46m 20.30s	+03° 02m 44.5s	0.66 ± 0.05	0.66 ± 0.01	0.64 ± 0.09	0.59 ± 0.06	0.57 ± 0.10	0.64 ± 0.08	1
NGC3110	10h 04m 02.11s	-06° 28m 29.6s	0.65 ± 0.03	0.66 ± 0.01	0.68 ± 0.03	0.66 ± 0.01	0.65 ± 0.03	0.68 ± 0.02	1
IRASF10038-3338	10h 06m 04.66s	-33° 53m 06.0s	0.05 ± 0.04	0.08 ± 0.03	0.39 ± 0.03	0.09 ± 0.09	0.06 ± 0.09	0.33 ± 0.04	3
IRASF10173+0828	10h 20m 00.24s	+08° 13m 32.9s	0.14 ± 0.09	0.14 ± 0.02	0.14 ± 0.01	0.11 ± 0.03	... ± ...	0.07 ± 0.09	0
NGC3221	10h 22m 19.97s	+21° 34m 10.6s	... ± ...	0.77 ± 0.02	... ± ...	0.80 ± 0.01	0.76 ± 0.04	... ± ...	1

Note. — Continued.

Table 2
FEEs of Various MIR Features

Galaxy name (1)	R.A. (J2000) (2)	Declination (J2000) (3)	FEE _{6.2μm} (PAH) (4)	FEE _{6.7μm} (Cont.) (5)	FEE _{7.7μm} (PAH) (6)	FEE _{9.7μm} (PAH) (7)	FEE _{11.3μm} (Si abs.) (8)	FEE _{12.8μm} ([Ne II]) (9)	FEE $_{\lambda}$ Type (10)
NGC3256	10h 27m 51.31s	-43° 54m 14.0s	0.69 ± 0.03	0.66 ± 0.01	0.47 ± 0.05	0.66 ± 0.01	0.57 ± 0.04	0.62 ± 0.03	2
ESO264-G036	10h 43m 07.51s	-46° 12m 44.3s	0.65 ± 0.02	0.65 ± 0.01	0.69 ± 0.02	0.65 ± 0.01	0.67 ± 0.03	0.70 ± 0.02	1
ESO264-G057	10h 59m 01.70s	-43° 26m 25.1s	0.33 ± 0.05	0.39 ± 0.02	0.44 ± 0.05	0.37 ± 0.06	0.38 ± 0.06	0.43 ± 0.04	1
IRASF10565+2448	10h 59m 18.14s	+24° 32m 34.1s	0.11 ± 0.07	0.15 ± 0.01	0.17 ± 0.05	0.19 ± 0.03	0.05 ± 0.08	0.16 ± 0.05	1
MCG+07-23-019	11h 03m 53.98s	+40° 51m 00.4s	0.50 ± 0.12	0.42 ± 0.21	0.57 ± 0.10	0.57 ± 0.09	0.62 ± 0.08	0.61 ± 0.09	4
CGCG011-076	11h 21m 12.24s	-02° 59m 02.4s	0.32 ± 0.03	0.32 ± 0.04	0.32 ± 0.04	0.41 ± 0.01	0.33 ± 0.06	0.39 ± 0.05	2
IC2810	11h 25m 45.07s	+14° 40m 36.1s	0.45 ± 0.03	0.49 ± 0.02	0.55 ± 0.03	0.51 ± 0.02	0.48 ± 0.04	0.56 ± 0.03	1
ESO319-G022	11h 27m 54.19s	-41° 36m 51.8s	0.17 ± 0.03	0.20 ± 0.07	0.18 ± 0.09	0.13 ± 0.05	0.10 ± 0.09	0.18 ± 0.06	1
ESO440-IG058	12h 06m 51.86s	-31° 56m 59.3s	0.56 ± 0.04	0.56 ± 0.01	0.61 ± 0.03	0.56 ± 0.01	0.54 ± 0.04	0.59 ± 0.03	1
ESO440-IG058	12h 06m 51.70s	-31° 56m 46.3s	0.52 ± 0.08	0.45 ± 0.09	0.50 ± 0.06	0.47 ± 0.01	0.50 ± 0.07	0.52 ± 0.03	1
IRASF12112+0305	12h 13m 46.03s	+02° 48m 42.1s	0.51 ± 0.08	0.46 ± 0.02	0.56 ± 0.05	0.49 ± 0.02	0.49 ± 0.06	0.52 ± 0.04	1
NGC4194	12h 14m 09.72s	+54° 31m 35.4s	0.48 ± 0.01	0.45 ± 0.04	0.50 ± 0.03	0.51 ± 0.01	0.47 ± 0.04	0.55 ± 0.04	1
ESO267-G030	12h 14m 12.82s	-47° 13m 42.6s	0.60 ± 0.02	0.60 ± 0.03	0.56 ± 0.03	0.61 ± 0.01	0.60 ± 0.03	0.62 ± 0.03	2
ESO267-G030	12h 13m 52.27s	-47° 16m 25.3s	0.33 ± 0.04	0.32 ± 0.04	0.36 ± 0.04	0.32 ± 0.05	0.27 ± 0.05	0.38 ± 0.03	1
IRAS12116-5615	12h 14m 22.08s	-56° 32m 32.6s	0.15 ± 0.07	0.27 ± 0.01	0.11 ± 0.06	0.25 ± 0.01	0.18 ± 0.05	0.28 ± 0.05	2
IRASF12224-0624	12h 25m 03.91s	-06° 40m 52.0s	0.20 ± 0.05	0.21 ± 0.10	0.37 ± 0.06	0.12 ± 0.08	0.10 ± 0.01	0.22 ± 0.05	3
Mrk231	12h 56m 14.26s	+56° 52m 25.0s	0.03 ± 0.01	0.18 ± 0.04	0.04 ± 0.01	0.13 ± 0.09	... ± ...	0.04 ± 0.01	0
NGC4922	13h 01m 25.27s	+29° 18m 49.7s	0.11 ± 0.04	0.13 ± 0.07	0.13 ± 0.06	0.11 ± 0.04	0.06 ± 0.08	0.10 ± 0.06	4
CGCG043-099	13h 01m 50.28s	+04° 20m 01.0s	0.13 ± 0.05	0.18 ± 0.03	0.37 ± 0.04	0.23 ± 0.07	0.10 ± 0.08	0.25 ± 0.05	3
MCG-02-33-098	13h 02m 19.66s	-15° 46m 04.1s	0.15 ± 0.09	0.19 ± 0.01	0.11 ± 0.04	0.25 ± 0.03	0.07 ± 0.08	0.20 ± 0.05	2
MCG-02-33-098	13h 02m 20.38s	-15° 45m 59.8s	0.49 ± 0.05	0.40 ± 0.06	0.45 ± 0.04	0.47 ± 0.02	0.43 ± 0.05	0.50 ± 0.04	1
ESO507-G070	13h 02m 52.42s	-23° 55m 17.8s	0.13 ± 0.07	0.17 ± 0.01	0.43 ± 0.04	0.15 ± 0.13	0.11 ± 0.08	0.22 ± 0.05	3
IRAS13052-5711	13h 08m 18.72s	-57° 27m 30.2s	0.37 ± 0.07	0.43 ± 0.03	0.52 ± 0.04	0.42 ± 0.01	0.40 ± 0.05	0.47 ± 0.05	3
IC0860	13h 15m 03.48s	+24° 37m 07.7s	0.29 ± 0.04	0.24 ± 0.06	0.51 ± 0.04	0.17 ± 0.08	0.21 ± 0.05	0.39 ± 0.03	3
IRAS13120-5453	13h 15m 06.36s	-55° 09m 22.3s	0.22 ± 0.04	0.17 ± 0.04	0.28 ± 0.05	0.23 ± 0.04	0.15 ± 0.06	0.26 ± 0.04	1
VV250a	13h 15m 34.97s	+62° 07m 29.3s	0.12 ± 0.06	0.14 ± 0.02	0.15 ± 0.04	0.16 ± 0.01	0.07 ± 0.06	0.17 ± 0.04	1
VV250a	13h 15m 30.70s	+62° 07m 45.8s	0.20 ± 0.08	0.24 ± 0.08	0.32 ± 0.13	0.27 ± 0.04	0.20 ± 0.06	0.28 ± 0.05	1
UGC08387	13h 20m 35.38s	+34° 08m 22.2s	0.17 ± 0.06	0.23 ± 0.01	0.36 ± 0.05	0.27 ± 0.01	0.16 ± 0.06	0.31 ± 0.05	3
NGC5104	13h 21m 23.09s	+00° 20m 33.4s	0.45 ± 0.04	0.45 ± 0.02	0.58 ± 0.03	0.46 ± 0.01	0.42 ± 0.06	0.53 ± 0.04	3
MCG-03-34-064	13h 22m 24.46s	-16° 43m 42.2s	0.02 ± 0.05	0.05 ± 0.02	0.05 ± 0.03	0.08 ± 0.03	0.04 ± 0.01	0.07 ± 0.05	0
NGC5135	13h 25m 44.02s	-29° 50m 00.2s	0.55 ± 0.03	0.54 ± 0.03	0.55 ± 0.03	0.55 ± 0.01	0.57 ± 0.04	0.59 ± 0.03	1
ESO173-G015	13h 27m 23.78s	-57° 29m 21.8s	0.35 ± 0.01	0.34 ± 0.05	0.56 ± 0.02	0.33 ± 0.07	0.29 ± 0.05	0.51 ± 0.04	3
IC4280	13h 32m 53.40s	-24° 12m 25.6s	0.76 ± 0.02	0.76 ± 0.01	0.78 ± 0.02	0.75 ± 0.01	0.76 ± 0.02	0.77 ± 0.02	1
NGC5256	13h 38m 17.26s	+48° 16m 32.9s	0.33 ± 0.04	0.39 ± 0.01	0.47 ± 0.04	0.39 ± 0.01	0.39 ± 0.06	0.42 ± 0.05	1
NGC5256	13h 38m 17.78s	+48° 16m 41.5s	0.49 ± 0.01	0.41 ± 0.01	0.44 ± 0.16	0.34 ± 0.10	0.29 ± 0.21	0.37 ± 0.19	1
NGC5257	13h 39m 57.72s	+00° 49m 53.0s	0.79 ± 0.03	0.78 ± 0.02	0.77 ± 0.03	0.77 ± 0.02	0.79 ± 0.03	0.78 ± 0.03	1
NGC5257	13h 39m 52.94s	+00° 50m 25.8s	0.76 ± 0.04	0.82 ± 0.03	0.83 ± 0.01	0.81 ± 0.03	0.82 ± 0.03	0.82 ± 0.02	1
Mrk273	13h 44m 42.12s	+55° 53m 13.2s	0.08 ± 0.03	0.24 ± 0.02	0.24 ± 0.08	0.18 ± 0.08	0.03 ± 0.08	0.18 ± 0.05	3
UGC08739	13h 49m 13.94s	+35° 15m 26.3s	0.37 ± 0.02	0.30 ± 0.07	0.61 ± 0.04	0.36 ± 0.07	0.45 ± 0.04	0.54 ± 0.02	3
ESO221-IG010	13h 50m 56.93s	-49° 03m 18.7s	0.44 ± 0.03	0.43 ± 0.04	0.45 ± 0.04	0.48 ± 0.01	0.43 ± 0.05	0.52 ± 0.04	1
NGC5331	13h 52m 16.20s	+02° 06m 05.0s	0.58 ± 0.03	0.57 ± 0.01	0.66 ± 0.02	0.58 ± 0.01	0.60 ± 0.03	0.64 ± 0.03	1
NGC5331	13h 52m 16.44s	+02° 06m 31.0s	0.59 ± 0.09	0.58 ± 0.07	0.65 ± 0.04	0.59 ± 0.04	0.62 ± 0.06	0.64 ± 0.05	1
NGC5395	13h 58m 37.97s	+37° 25m 28.2s	0.62 ± 0.12	0.67 ± 0.07	0.63 ± 0.04	0.71 ± 0.01	0.71 ± 0.01	0.71 ± 0.01	1
NGC5395	13h 58m 33.65s	+37° 27m 13.0s	0.31 ± 0.03	0.31 ± 0.02	0.33 ± 0.03	0.35 ± 0.01	0.30 ± 0.05	0.34 ± 0.04	1
CGCG247-020	14h 19m 43.27s	+49° 14m 11.8s	0.03 ± 0.07	0.14 ± 0.01	0.11 ± 0.06	0.19 ± 0.05	0.06 ± 0.08	0.12 ± 0.06	1
NGC5653	14h 30m 10.44s	+31° 12m 55.8s	0.75 ± 0.01	0.76 ± 0.02	0.76 ± 0.02	0.76 ± 0.01	0.76 ± 0.02	0.77 ± 0.02	1
IRASF14348-1447*	14h 37m 38.28s	-15° 00m 24.1s	0.41 ± 0.03	0.50 ± 0.04	0.58 ± 0.03	0.53 ± 0.03	0.56 ± 0.05	0.60 ± 0.03	4
IRASF14378-3651	14h 40m 59.04s	-37° 04m 32.2s	0.16 ± 0.07	0.11 ± 0.07	0.29 ± 0.12	0.13 ± 0.03	0.05 ± 0.01	0.08 ± 0.07	4
NGC5734	14h 45m 09.05s	-20° 52m 13.1s	0.77 ± 0.01	0.79 ± 0.02	0.77 ± 0.02	0.78 ± 0.01	0.77 ± 0.03	0.78 ± 0.03	1

Note. — Continued.

Table 2
FEEs of Various MIR Features

Galaxy name (1)	R.A. (J2000) (2)	Declination (J2000) (3)	FEE _{6.2μm} (PAH) (4)	FEE _{6.7μm} (Cont.) (5)	FEE _{7.7μm} (PAH) (6)	FEE _{9.7μm} (PAH) (7)	FEE _{11.3μm} (Si abs.) (8)	FEE _{12.8μm} ([Ne II]) (9)	FEE _{λ} Type (10)
NGC5734	14h 45m 11.02s	-20° 54m 48.6s	0.67 ± 0.02	0.70 ± 0.02	0.68 ± 0.06	0.67 ± 0.04	0.68 ± 0.06	0.69 ± 0.05	1
VV340a	14h 57m 00.70s	+24° 37m 05.9s	0.78 ± 0.02	0.79 ± 0.01	0.81 ± 0.02	0.78 ± 0.01	0.77 ± 0.02	0.80 ± 0.02	1
VV340a	14h 57m 00.31s	+24° 36m 24.5s	0.70 ± 0.04	0.67 ± 0.03	0.63 ± 0.06	0.61 ± 0.05	0.64 ± 0.08	0.65 ± 0.07	1
VV705	15h 18m 06.14s	+42° 44m 44.9s	0.52 ± 0.23	0.41 ± 0.07	0.65 ± 0.10	0.41 ± 0.02	0.52 ± 0.10	0.50 ± 0.12	4
ESO099-G004	15h 24m 57.98s	-63° 07m 29.3s	0.50 ± 0.05	0.55 ± 0.07	0.41 ± 0.03	0.48 ± 0.02	0.24 ± 0.08	0.46 ± 0.04	2
IRASF15250+3608	15h 26m 59.42s	+35° 58m 37.9s	0.05 ± 0.03	0.05 ± 0.04	0.18 ± 0.10	0.08 ± 0.06	... ± ...	0.17 ± 0.05	3
NGC5936	15h 30m 00.84s	+12° 59m 22.2s	0.39 ± 0.01	0.32 ± 0.06	0.39 ± 0.05	0.38 ± 0.07	0.35 ± 0.04	0.42 ± 0.02	1
Arp220	15h 34m 57.24s	+23° 30m 11.2s	0.19 ± 0.06	0.24 ± 0.01	0.48 ± 0.04	0.16 ± 0.08	0.10 ± 0.09	0.43 ± 0.03	3
NGC5990	15h 46m 16.42s	+02° 24m 55.4s	0.27 ± 0.05	0.34 ± 0.01	0.30 ± 0.05	0.44 ± 0.02	0.36 ± 0.05	0.48 ± 0.04	2
NGC6090	16h 11m 40.85s	+52° 27m 27.4s	0.50 ± 0.03	0.50 ± 0.02	0.53 ± 0.03	0.54 ± 0.01	0.53 ± 0.04	0.55 ± 0.03	1
IRASF16164-0746	16h 19m 11.76s	-07° 54m 02.9s	0.17 ± 0.06	0.20 ± 0.02	0.52 ± 0.05	0.25 ± 0.03	0.12 ± 0.08	0.37 ± 0.05	3
CGCG052-037	16h 30m 53.26s	+04° 04m 23.9s	0.35 ± 0.03	0.35 ± 0.04	0.41 ± 0.03	0.38 ± 0.04	0.35 ± 0.04	0.41 ± 0.03	1
NGC6156	16h 34m 52.56s	-60° 37m 08.0s	0.41 ± 0.02	0.46 ± 0.03	0.14 ± 0.04	0.46 ± 0.01	0.23 ± 0.06	0.32 ± 0.04	2
ESO069-IG006	16h 38m 11.86s	-68° 26m 08.2s	0.39 ± 0.07	0.40 ± 0.03	0.51 ± 0.04	0.42 ± 0.01	0.41 ± 0.06	0.46 ± 0.04	1
IRASF16399-0937*	16h 42m 40.10s	-09° 43m 13.8s	0.72 ± 0.02	0.73 ± 0.01	0.74 ± 0.02	0.71 ± 0.01	0.61 ± 0.04	0.73 ± 0.02	4
ESO453-G005	16h 47m 31.08s	-29° 21m 21.6s	0.70 ± 0.02	0.73 ± 0.01	0.73 ± 0.01	0.70 ± 0.02	0.69 ± 0.03	0.71 ± 0.02	1
NGC6240*	16h 52m 58.90s	+02° 24m 03.2s	0.32 ± 0.06	0.34 ± 0.01	0.48 ± 0.04	0.35 ± 0.02	0.31 ± 0.05	0.45 ± 0.04	3
IRASF16516-0948	16h 54m 23.71s	-09° 53m 20.8s	0.71 ± 0.02	0.72 ± 0.02	0.75 ± 0.04	0.71 ± 0.03	0.73 ± 0.05	0.74 ± 0.04	1
NGC6286	16h 58m 31.63s	+58° 56m 13.2s	0.80 ± 0.04	0.80 ± 0.03	0.80 ± 0.01	0.80 ± 0.01	0.80 ± 0.02	0.80 ± 0.01	1
NGC6286	16h 58m 24.00s	+58° 57m 21.6s	0.50 ± 0.10	0.47 ± 0.01	0.59 ± 0.03	0.48 ± 0.02	0.49 ± 0.05	0.55 ± 0.03	3
IRASF17132+5313*	17h 14m 20.45s	+53° 10m 31.4s	0.66 ± 0.03	0.67 ± 0.02	0.68 ± 0.03	0.66 ± 0.01	0.70 ± 0.03	0.70 ± 0.03	1
IRASF17138-1017	17h 16m 35.69s	-10° 20m 40.6s	0.65 ± 0.03	0.64 ± 0.03	0.69 ± 0.02	0.66 ± 0.01	0.70 ± 0.03	0.71 ± 0.03	1
IRASF17207-0014	17h 23m 21.96s	+00° 17m 00.6s	0.17 ± 0.06	0.24 ± 0.01	0.45 ± 0.04	0.25 ± 0.03	0.19 ± 0.05	0.40 ± 0.04	3
ESO138-G027	17h 26m 43.34s	-59° 55m 55.2s	0.48 ± 0.03	0.55 ± 0.03	0.36 ± 0.04	0.55 ± 0.01	0.36 ± 0.06	0.54 ± 0.04	2
UGC11041	17h 54m 51.82s	+34° 46m 34.3s	0.56 ± 0.02	0.56 ± 0.03	0.61 ± 0.02	0.57 ± 0.02	0.60 ± 0.03	0.62 ± 0.02	1
CGCG141-034	17h 56m 56.64s	+24° 01m 01.9s	0.23 ± 0.05	0.25 ± 0.02	0.42 ± 0.06	0.20 ± 0.05	0.20 ± 0.07	0.30 ± 0.04	3
IRAS17578-0400	18h 00m 31.85s	-04° 00m 53.3s	0.39 ± 0.06	0.40 ± 0.07	0.55 ± 0.04	0.44 ± 0.01	0.48 ± 0.05	0.54 ± 0.03	4
IRAS17578-0400	18h 00m 34.08s	-04° 01m 44.0s	0.43 ± 0.05	0.49 ± 0.02	0.54 ± 0.04	0.50 ± 0.01	0.50 ± 0.04	0.56 ± 0.03	1
IRAS17578-0400	18h 00m 24.29s	-04° 01m 03.7s	0.60 ± 0.05	0.60 ± 0.02	0.68 ± 0.04	0.64 ± 0.01	0.67 ± 0.03	0.68 ± 0.03	1
IRAS18090+0130	18h 11m 38.42s	+01° 31m 40.4s	0.52 ± 0.05	0.55 ± 0.02	0.63 ± 0.04	0.55 ± 0.01	0.51 ± 0.04	0.59 ± 0.03	1
IRAS18090+0130	18h 11m 33.41s	+01° 31m 42.2s	0.35 ± 0.07	0.34 ± 0.03	0.54 ± 0.09	0.38 ± 0.06	0.32 ± 0.05	0.46 ± 0.04	3
CGCG142-034	18h 16m 40.68s	+22° 06m 46.4s	0.24 ± 0.03	0.23 ± 0.04	0.46 ± 0.03	0.21 ± 0.11	0.33 ± 0.04	0.40 ± 0.03	3
CGCG142-034	18h 16m 33.84s	+22° 06m 38.9s	0.47 ± 0.04	0.49 ± 0.02	0.55 ± 0.05	0.47 ± 0.01	0.44 ± 0.05	0.49 ± 0.04	1
IRASF18293-3413	18h 32m 41.09s	-34° 11m 26.9s	0.45 ± 0.04	0.49 ± 0.01	0.54 ± 0.03	0.51 ± 0.01	0.51 ± 0.04	0.54 ± 0.03	1
NGC6670AB	18h 33m 34.25s	+59° 53m 17.9s	0.25 ± 0.06	0.27 ± 0.01	0.40 ± 0.03	0.26 ± 0.05	0.32 ± 0.06	0.39 ± 0.04	3
NGC6670AB	18h 33m 37.73s	+59° 53m 22.9s	0.24 ± 0.06	0.25 ± 0.01	0.32 ± 0.03	0.28 ± 0.05	0.23 ± 0.07	0.35 ± 0.03	1
IC4734	18h 38m 25.75s	-57° 29m 25.4s	0.21 ± 0.05	0.23 ± 0.01	0.40 ± 0.04	0.30 ± 0.06	0.22 ± 0.07	0.35 ± 0.04	3
NGC6701	18h 43m 12.53s	+60° 39m 11.5s	0.41 ± 0.05	0.46 ± 0.01	0.48 ± 0.03	0.47 ± 0.02	0.45 ± 0.05	0.48 ± 0.04	1
NGC6786	19h 10m 54.00s	+73° 24m 36.0s	0.54 ± 0.04	0.53 ± 0.02	0.56 ± 0.04	0.54 ± 0.01	0.53 ± 0.04	0.59 ± 0.03	1
NGC6786	19h 11m 04.37s	+73° 25m 32.5s	0.36 ± 0.05	0.47 ± 0.04	0.24 ± 0.08	0.45 ± 0.04	0.21 ± 0.08	0.34 ± 0.04	2
ESO593-IG008	19h 14m 31.15s	-21° 19m 06.2s	0.56 ± 0.05	0.58 ± 0.01	0.63 ± 0.03	0.53 ± 0.01	0.55 ± 0.05	0.61 ± 0.03	1
IRASF19297-0406	19h 32m 22.30s	-04° 00m 01.1s	0.16 ± 0.04	0.18 ± 0.04	0.25 ± 0.12	0.15 ± 0.04	0.13 ± 0.08	0.18 ± 0.06	1
IRAS19542+1110	19h 56m 35.78s	+11° 19m 04.8s	0.11 ± 0.03	0.09 ± 0.06	0.10 ± 0.07	0.06 ± 0.03	0.03 ± 0.08	0.05 ± 0.06	0
ESO339-G011	19h 57m 37.61s	-37° 56m 08.5s	0.29 ± 0.03	0.36 ± 0.04	0.25 ± 0.04	0.38 ± 0.04	0.23 ± 0.06	0.38 ± 0.03	2
NGC6907	20h 25m 06.58s	-24° 48m 32.8s	0.64 ± 0.04	0.62 ± 0.01	0.64 ± 0.02	0.64 ± 0.01	0.60 ± 0.04	0.66 ± 0.03	1
MCG+04-48-002	20h 28m 35.06s	+25° 44m 00.2s	0.41 ± 0.03	0.43 ± 0.04	0.52 ± 0.03	0.46 ± 0.01	0.48 ± 0.04	0.52 ± 0.04	1
NGC6926	20h 33m 06.12s	-02° 01m 39.0s	0.73 ± 0.08	0.85 ± 0.01	0.85 ± 0.01	0.85 ± 0.01	0.79 ± 0.02	0.85 ± 0.01	4
IRAS20351+2521	20h 37m 17.74s	+25° 31m 37.6s	0.40 ± 0.07	0.45 ± 0.02	0.54 ± 0.04	0.53 ± 0.02	0.44 ± 0.06	0.56 ± 0.04	1
CGCG448-020	20h 57m 24.10s	+17° 07m 35.0s	0.49 ± 0.08	0.49 ± 0.02	0.52 ± 0.05	0.55 ± 0.02	0.60 ± 0.05	0.56 ± 0.03	1

Note. — Continued.

Table 2
FEEs of Various MIR Features

Galaxy name (1)	R.A. (J2000) (2)	Declination (J2000) (3)	FEE _{6.2μm} (PAH) (4)	FEE _{6.7μm} (Cont.) (5)	FEE _{7.7μm} (PAH) (6)	FEE _{9.7μm} (PAH) (7)	FEE _{11.3μm} (Si abs.) (8)	FEE _{12.8μm} ([Ne II]) (9)	FEE $_{\lambda}$ Type (10)
CGCG448-020	20h 57m 24.38s	+17° 07m 39.4s	0.13 ± 0.09	0.20 ± 0.02	0.20 ± 0.06	0.23 ± 0.03	0.13 ± 0.06	0.31 ± 0.04	2
ESO286-IG019	20h 58m 26.78s	-42° 39m 00.4s	0.12 ± 0.09	0.20 ± 0.04	0.41 ± 0.04	0.22 ± 0.08	0.05 ± 0.08	0.26 ± 0.05	3
ESO286-G035	21h 04m 11.11s	-43° 35m 36.2s	0.57 ± 0.02	0.56 ± 0.02	0.62 ± 0.02	0.56 ± 0.01	0.59 ± 0.03	0.61 ± 0.03	1
IRAS21101+5810	21h 11m 29.28s	+58° 23m 07.8s	0.27 ± 0.07	0.26 ± 0.03	0.25 ± 0.08	0.29 ± 0.02	0.11 ± 0.08	0.25 ± 0.06	4
ESO343-IG013	21h 36m 10.54s	-38° 32m 42.7s	0.40 ± 0.06	0.41 ± 0.01	0.61 ± 0.03	0.42 ± 0.01	0.47 ± 0.04	0.53 ± 0.04	3
ESO343-IG013	21h 36m 10.92s	-38° 32m 33.0s	0.18 ± 0.06	0.28 ± 0.02	0.19 ± 0.05	0.25 ± 0.06	0.04 ± 0.06	0.25 ± 0.05	2
NGC7130	21h 48m 19.54s	-34° 57m 04.7s	0.46 ± 0.03	0.53 ± 0.05	0.47 ± 0.07	0.57 ± 0.05	0.48 ± 0.08	0.59 ± 0.06	2
ESO467-G027	22h 14m 39.96s	-27° 27m 50.4s	0.71 ± 0.02	0.73 ± 0.02	0.73 ± 0.02	0.72 ± 0.01	0.72 ± 0.02	0.74 ± 0.02	1
ESO602-G025	22h 31m 25.49s	-19° 02m 03.8s	0.35 ± 0.06	0.44 ± 0.01	0.52 ± 0.04	0.43 ± 0.01	0.39 ± 0.05	0.47 ± 0.04	3
UGC12150	22h 41m 12.22s	+34° 14m 56.8s	0.33 ± 0.06	0.39 ± 0.02	0.43 ± 0.04	0.39 ± 0.03	0.35 ± 0.04	0.43 ± 0.04	1
IRASF22491-1808	22h 51m 49.34s	-17° 52m 25.0s	0.19 ± 0.03	0.27 ± 0.06	0.35 ± 0.07	0.35 ± 0.02	0.27 ± 0.08	0.39 ± 0.06	2
NGC7469	23h 03m 15.65s	+08° 52m 25.7s	0.25 ± 0.06	0.37 ± 0.01	0.15 ± 0.06	0.41 ± 0.01	0.30 ± 0.05	0.35 ± 0.04	2
CGCG453-062	23h 04m 56.54s	+19° 33m 07.2s	0.69 ± 0.02	0.72 ± 0.03	0.77 ± 0.03	0.72 ± 0.02	0.71 ± 0.03	0.75 ± 0.02	3
ESO148-IG002	23h 15m 46.75s	-59° 03m 15.8s	0.13 ± 0.02	0.35 ± 0.05	0.17 ± 0.10	0.31 ± 0.04	0.06 ± 0.09	0.27 ± 0.06	2
IC5298	23h 16m 00.67s	+25° 33m 24.5s	0.15 ± 0.05	0.26 ± 0.04	0.07 ± 0.04	0.20 ± 0.02	0.09 ± 0.05	0.18 ± 0.05	2
NGC7591	23h 18m 16.25s	+06° 35m 09.2s	0.44 ± 0.05	0.39 ± 0.04	0.49 ± 0.04	0.38 ± 0.03	0.31 ± 0.04	0.45 ± 0.03	3
ESO077-IG014	23h 21m 05.45s	-69° 12m 47.2s	0.14 ± 0.09	0.16 ± 0.01	0.32 ± 0.04	0.20 ± 0.05	0.09 ± 0.09	0.21 ± 0.04	3
ESO077-IG014	23h 21m 03.72s	-69° 13m 00.8s	0.18 ± 0.07	0.20 ± 0.02	0.37 ± 0.05	0.24 ± 0.03	0.15 ± 0.09	0.26 ± 0.05	3
IRASF23365+3604	23h 39m 01.32s	+36° 21m 08.3s	0.37 ± 0.08	0.28 ± 0.03	0.49 ± 0.17	0.38 ± 0.02	0.16 ± 0.12	0.43 ± 0.06	3
MCG-01-60-022	23h 42m 00.91s	-03° 36m 54.4s	0.43 ± 0.07	0.48 ± 0.01	0.60 ± 0.04	0.47 ± 0.02	0.42 ± 0.06	0.59 ± 0.04	3
IRAS23436+5257	23h 46m 05.45s	+53° 14m 01.7s	0.35 ± 0.05	0.39 ± 0.05	0.31 ± 0.04	0.45 ± 0.04	0.36 ± 0.08	0.41 ± 0.05	2
NGC7753	23h 47m 04.85s	+29° 29m 00.6s	0.39 ± 0.06	0.37 ± 0.03	0.41 ± 0.06	0.37 ± 0.02	0.32 ± 0.04	0.41 ± 0.04	1
NGC7753	23h 46m 58.63s	+29° 27m 32.0s	0.62 ± 0.03	0.62 ± 0.02	0.67 ± 0.02	0.63 ± 0.01	0.63 ± 0.03	0.68 ± 0.03	1
NGC7771	23h 51m 03.91s	+20° 09m 01.8s	0.55 ± 0.07	0.56 ± 0.02	0.51 ± 0.04	0.57 ± 0.05	0.58 ± 0.05	0.65 ± 0.05	2
NGC7771	23h 51m 24.79s	+20° 06m 42.1s	0.46 ± 0.01	0.41 ± 0.05	0.52 ± 0.03	0.48 ± 0.01	0.50 ± 0.04	0.53 ± 0.03	1
NGC7771	23h 51m 22.56s	+20° 05m 49.2s	0.48 ± 0.05	0.56 ± 0.05	0.52 ± 0.04	0.59 ± 0.01	0.58 ± 0.04	0.64 ± 0.03	2
Mrk0331	23h 51m 26.76s	+20° 35m 10.3s	0.38 ± 0.03	0.40 ± 0.04	0.46 ± 0.04	0.43 ± 0.01	0.38 ± 0.05	0.44 ± 0.04	1

Note. — Continued.